

THE TWO OBSTACLE PROBLEM FOR THE PARABOLIC BIHARMONIC EQUATION

M. NOVAGA AND S. OKABE

ABSTRACT. We consider a two obstacle problem for the parabolic biharmonic equation in a bounded domain. We prove long time existence of solutions via an implicit time discretization scheme, and we investigate the regularity properties of solutions.

1. INTRODUCTION

The present paper is devoted to discussing a two obstacle problem for the parabolic biharmonic equation. The obstacle problem for second order elliptic and parabolic equations has attracted a great interest in the past years, and there is an extensive mathematical literature (e.g., see [6] and the references therein). On the contrary, much less is known on the obstacle problem for higher order elliptic or parabolic equations.

The biharmonic operator can be regarded as a prototype fourth order differential operator. Indeed, elliptic and parabolic PDEs for biharmonic operator are under intensive investigation in recent years (see for example [2, 9, 10, 12, 13, 14, 15, 16]). Although the obstacle problem for the biharmonic equation has been studied in the 1970s and 1980s (see [?, 5, 7, 8, 11, 20]), some results on the obstacle problem for the corresponding parabolic equation have only been obtained very recently. In particular, in [19] we considered the case of a single obstacle, i.e., the solution u satisfies $u \geq f$ in Ω for a given obstacle function f in a domain Ω , and it is natural to ask whether the results can be extended to the case of two obstacles. Indeed, in this paper we prove the existence of solutions for the two obstacle problem, and we investigate their regularity properties.

Let $\Omega \subset \mathbb{R}^N$, with $N \leq 3$, be a bounded domain with $\partial\Omega \in C^4$. Let $f : \Omega \rightarrow \mathbb{R}$ and $g : \Omega \rightarrow \mathbb{R}$ denote the obstacle functions satisfying

$$(1.1) \quad f \in C^4(\overline{\Omega}), \quad g \in C^4(\overline{\Omega}), \quad f \leq g \quad \text{in } \Omega,$$

$$(1.2) \quad f < 0 < g \quad \text{on } \partial\Omega.$$

We consider a two obstacle problem of the type

$$(P) \quad \begin{cases} (\partial_t u + \Delta^2 u)(u - f) \leq 0 & \text{in } \Omega \times \mathbb{R}_+, \\ (\partial_t u + \Delta^2 u)(u - g) \leq 0 & \text{in } \Omega \times \mathbb{R}_+, \\ \partial_t u + \Delta^2 u = 0 & \text{in } \{(x, t) \in \Omega \times \mathbb{R}_+ \mid f(x) < u(x, t) < g(x)\}, \\ f \leq u \leq g & \text{in } \Omega \times \mathbb{R}_+, \\ u = \nabla u \cdot \nu^\Omega = 0 & \text{on } \partial\Omega \times \mathbb{R}_+, \\ u(\cdot, 0) = u_0(\cdot) & \text{in } \Omega, \end{cases}$$

1991 *Mathematics Subject Classification.* 35K25, 35J35, 49A29.

Key words and phrases. minimizing movements.

The second author was partially supported by Grant-in-Aid for Young Scientists (B), No. 24740097, and by Strategic Young Researcher Overseas Visits Program for Accelerating Brain Circulation (JSPS).

where ν^Ω denotes the unit normal vector on $\partial\Omega$, and the initial datum $u_0 : \Omega \rightarrow \mathbb{R}$ satisfies

$$(1.3) \quad u_0 \in H_0^2(\Omega), \quad f \leq u_0 \leq g \quad \text{in } \Omega.$$

Here we define a weak solution of (P). To this aim, we set

$$(1.4) \quad \mathcal{K} := \{ u \in L^2(0, T; H_0^2(\Omega)) \cap H^1(0, T; L^2(\Omega)) \mid u(x, 0) = u_0(x) \text{ a.e. in } \Omega, \\ f(x) \leq u(x, t) \leq g(x) \text{ a.e. in } \Omega \times (0, T) \}.$$

Definition 1.1. *We say that a function u is a weak solution of (P) if*

- (i) $u \in \mathcal{K}$;
- (ii) for any $w \in \mathcal{K}$,

$$(1.5) \quad \int_0^T \int_\Omega [\partial_t u(w - u) + \Delta u \Delta(w - u)] \, dx dt \geq 0.$$

Let us denote by Ω_0 the coincidence set of f and g , i.e.,

$$(1.6) \quad \Omega_0 = \{ x \in \Omega \mid f(x) = g(x) \}.$$

The main result of this paper is the following:

Theorem 1.1. *Let $N \leq 3$. Let f and g satisfy (1.1)-(1.2). Then, for any initial datum u_0 satisfying (1.3), the problem (P) possesses a unique weak solution*

$$(1.7) \quad u \in L^\infty(\mathbb{R}_+; H_0^2(\Omega)) \cap H^1(\mathbb{R}_+; L^2(\Omega)).$$

Moreover the quantity $\mu_t := \partial_t u(\cdot, t) + \Delta^2 u(\cdot, t)$ defines a signed measure in Ω for a.e. $t \in \mathbb{R}_+$, and for any $T > 0$ there exists a constant $C > 0$ such that

$$(1.8) \quad \int_0^T \mu_t(\Omega)^2 \, dt < C + T \|\Delta^2 f\|_{L^\infty(\Omega_0)}^2.$$

Furthermore the following regularity properties hold:

- (i) $u \in L^2(\mathbb{R}_+; W^{2,\infty}(\Omega))$. In particular, if $N = 1$,

$$(1.9) \quad u \in C^{0,\beta}(\mathbb{R}_+; C^{1,\gamma}(\Omega)) \quad \text{with } 0 < \gamma < \frac{1}{2} \text{ and } 0 < \beta < \frac{1-2\gamma}{8},$$

if $N \in \{2, 3\}$,

$$(1.10) \quad u \in C^{0,\beta}(\mathbb{R}_+; C^{0,\gamma}(\Omega)) \quad \text{with } 0 < \gamma < \frac{4-N}{2} \text{ and } 0 < \beta < \frac{4-N-2\gamma}{8};$$

- (ii) the signed measure μ_t satisfies

$$(1.11) \quad \mu_t \llcorner_{\Omega_0} = \Delta^2 f,$$

$$(1.12) \quad \text{supp } \mu_t \llcorner_{\Omega \setminus \Omega_0} \subset \{ (x, t) \in (\Omega \setminus \Omega_0) \times \mathbb{R}_+ \mid u(x, t) = f(x) \text{ or } u(x, t) = g(x) \},$$

with

$$(1.13) \quad \mu_t \begin{cases} \geq 0 & \text{in } \{ (x, t) \in (\Omega \setminus \Omega_0) \times \mathbb{R}_+ \mid u(x, t) = f(x) \}, \\ \leq 0 & \text{in } \{ (x, t) \in (\Omega \setminus \Omega_0) \times \mathbb{R}_+ \mid u(x, t) = g(x) \}. \end{cases}$$

In particular, u satisfies (P) in the sense of distributions.

The restriction on the dimension $N \leq 3$ in Theorem 1.1 has two motivations. The first is related to the continuity of the approximate solutions. We construct the solution of (P) as a suitable limit of solutions of the obstacle problem for the corresponding elliptic equation, which is a biharmonic equation with a lower order perturbation. Here a difficulty arises from the presence of the set Ω_0 . To overcome this difficulty, first we construct the solution of the two obstacle problem replaced f with $f - \varepsilon$, for $\varepsilon > 0$. If the solution u_ε of the modified two obstacle problem is uniformly continuous with respect to ε in Ω , then one can obtain a solution of the original obstacle problem as a limit of u_ε as $\varepsilon \downarrow 0$. Thus the point is to obtain the uniform continuity of u_ε , and this is given by Sobolev's embedding if $N \leq 3$. For the same reason, the two obstacle problem for the elliptic biharmonic equation was studied in [8] under the same assumption $N \leq 3$.

Even if $\Omega_0 = \emptyset$, we still need the restriction on the dimension in order to prove the $C^{1,1}$ regularity of the approximate solutions. Here the difficulty proving the continuity of the discrete velocities, which converge to $\partial_t u$. Again, such continuity can be obtained from Sobolev's embedding if $N \leq 3$.

We note that Theorem 1.1 can be extended to the problem (P) replaced Neumann boundary condition by Navier boundary condition, i.e., $u = \Delta u = 0$ on $\partial\Omega$. Indeed, replacing $H_0^2(\Omega)$ by $H^2(\Omega) \cap H_0^1(\Omega)$, we obtain the same conclusion as Theorem 1.1

The paper is organized as follows: We shall construct the solution of (P) by way of an implicit time discretization so called minimizing movements, which was given by De Giorgi. We give a formulation via minimizing movement in Section 2. In Section 3, we construct an approximate solution of the problem (P) and investigate its regularity. In Section 4, we prove Theorem 1.1. Indeed, we first prove that the approximate solution converges to a function in a suitable sense. And then we observe that the limit is the required solution of (P).

2. NOTATION

In this paper, we shall construct a solution of (P) via minimizing movements (i.e., see [1]). We first note that the problem (P) is the L^2 -gradient flow for the functional

$$(2.1) \quad E(u) := \frac{1}{2} \int_{\Omega} |\Delta u(x)|^2 dx$$

with constraint $u \in \mathcal{K}$. Let $T > 0$ and $n \in \mathbb{N}$, and set $\tau_n = T/n$. We define a sequence $\{u_{i,n}\}_{i=0}^n$ inductively. To begin with, we let $u_{0,n} := u_0$. Let us denote by $u_{i,n}$ the minimizer of the problem

$$(M_{i,n}) \quad \min\{G_{i,n}(u) \mid u \in K\}$$

with

$$(2.2) \quad G_{i,n}(u) := E(u) + P_{i,n}(u)$$

where

$$(2.3) \quad P_{i,n}(u) := \frac{1}{2\tau_n} \int_{\Omega} [u(x) - u_{i-1,n}(x)]^2 dx.$$

The set K is given by

$$(2.4) \quad K = \{u \in H_0^2(\Omega) \mid f \leq u \leq g \text{ in } \Omega\}.$$

Let us set

$$(2.5) \quad V_{i,n}(x) = \frac{u_{i,n}(x) - u_{i-1,n}(x)}{\tau_n}.$$

Definition 2.1. Let us define $u_n(x, t) : \Omega \times [0, T] \rightarrow \mathbb{R}$ as

$$(2.6) \quad u_n(x, t) = u_{i-1,n}(x) + (t - (i-1)\tau_n)V_{i,n}(x)$$

in $\Omega \times [(i-1)\tau_n, i\tau_n]$ for each $i = 1, 2, \dots, n$.

Definition 2.2. Let us define $\tilde{u}_n(x, t) : \Omega \times (0, T] \rightarrow \mathbb{R}$ and $V_n(x, t) : \Omega \times (0, T] \rightarrow \mathbb{R}$ as

$$(2.7) \quad \tilde{u}_n(x, t) = u_{i,n}(x),$$

$$(2.8) \quad V_n(x, t) = V_{i,n}(x),$$

in $\Omega \times ((i-1)\tau_n, i\tau_n]$ for each $i = 1, 2, \dots, n$.

3. EXISTENCE OF APPROXIMATE SOLUTION

To begin with, we show the existence of the solution of $(M_{i,n})$.

Theorem 3.1. Let f and g satisfy (1.1)-(1.2). Let u_0 satisfy (1.3). Then there exists a unique minimizer of $(M_{i,n})$.

Proof. Let $\{u_j\} \subset K$ be a minimizing sequence for the functional (2.2). Since

$$0 \leq \inf_K G_{i,n}(u) \leq G_{i,n}(u_{i-1,n}) = E(u_{i-1,n}),$$

we may assume $\{u_j\}$ that $\sup_{j \in \mathbb{N}} G_{i,n}(u_j) < \infty$. Recalling that $\|\Delta v\|_{L^2(\Omega)}$ is equivalent to $\|v\|_{H_0^2(\Omega)}$ on $H_0^2(\Omega)$, we deduce that $\{u_j\}$ is uniformly bounded in $H_0^2(\Omega)$, and then there exists $u \in H_0^2(\Omega)$ such that

$$(3.1) \quad u_j \rightharpoonup u \quad \text{in} \quad H^2(\Omega),$$

in particular,

$$(3.2) \quad \Delta u_j \rightharpoonup \Delta u \quad \text{in} \quad L^2(\Omega),$$

up to a subsequence. Since (3.1) implies that u_j uniformly converges to u in Ω up to a subsequence, we have $f \leq u \leq g$ in Ω . It follows from Fatou's Lemma that

$$P_{i,n}(u) \leq \liminf_{j \rightarrow \infty} P_{i,n}(u_j).$$

Moreover we infer from (3.2) that

$$E(u) \leq \liminf_{j \rightarrow \infty} E(u_j).$$

The uniqueness of the minimizer of $(M_{i,n})$ follows from the convexity of $G_{i,n}$. \square

Set

$$(3.3) \quad f_\varepsilon(x) = f(x) - \varepsilon.$$

We denote by $(M_{i,n}^\varepsilon)$ the problem $(M_{i,n})$ replaced f by f_ε . The proof of Theorem 3.1 implies that the problem $(M_{i,n}^\varepsilon)$ has a unique minimizer $u_{i,n}^\varepsilon$. From now on, let us set

$$(3.4) \quad V_{i,n}^\varepsilon = \frac{u_{i,n}^\varepsilon - u_{i-1,n}^\varepsilon}{\tau_n}.$$

Moreover let V_n^ε denote the piecewise constant interpolation of $V_{i,n}^\varepsilon$.

Lemma 3.1. $u_{i,n}^\varepsilon$ uniformly converges to $u_{i,n}$ in Ω as $\varepsilon \rightarrow 0$.

Proof. By the fact that $\|u_{i,n}^\varepsilon\|_{H^2(\Omega)} \leq C$, for any sequence $\{\varepsilon_m\}$ with $\varepsilon_m \rightarrow 0$ as $m \rightarrow \infty$, there exist $\{\varepsilon_{m'}\} \subset \{\varepsilon_m\}$ and $\bar{u}_{i,n} \in H_0^2(\Omega)$ such that

$$(3.5) \quad u_{i,n}^{\varepsilon_{m'}} \rightharpoonup \bar{u}_{i,n} \quad \text{weakly in } H^2(\Omega) \quad \text{as } m' \rightarrow \infty,$$

in particular,

$$(3.6) \quad \Delta u_{i,n}^{\varepsilon_{m'}} \rightharpoonup \Delta \bar{u}_{i,n} \quad \text{weakly in } L^2(\Omega) \quad \text{as } m' \rightarrow \infty.$$

Since $N \leq 3$, Sobolev's embedding theorem implies that $u_{i,n}^{\varepsilon_{m'}}$ uniformly converges to $\bar{u}_{i,n}$ as $\varepsilon \downarrow 0$. Recalling that the solution $u_{i,n}^{\varepsilon_{m'}}$ of $(M_{i,n}^{\varepsilon_{m'}})$ satisfies

$$\int_{\Omega} [\Delta u_{i,n}^{\varepsilon_{m'}} \Delta(w - u_{i,n}^{\varepsilon_{m'}}) + V_{i,n}^{\varepsilon_{m'}}(w - u_{i,n}^{\varepsilon_{m'}})] dx \geq 0 \quad \text{for any } w \in K_{\varepsilon_{m'}},$$

we deduce from (3.5)-(3.6) that

$$\begin{aligned} & \int_{\Omega} [\Delta \bar{u}_{i,n} \Delta(w - \bar{u}_{i,n}) + \bar{V}_{i,n}(w - \bar{u}_{i,n})] dx \\ & \geq \liminf_{m' \rightarrow \infty} \int_{\Omega} [\Delta u_{i,n}^{\varepsilon_{m'}} \Delta(w - u_{i,n}^{\varepsilon_{m'}}) + V_{i,n}^{\varepsilon_{m'}}(w - u_{i,n}^{\varepsilon_{m'}})] dx \geq 0 \quad \text{for any } w \in K, \end{aligned}$$

where we used the fact $K \subset K_{\varepsilon_{m'}}$. Moreover it follows from the uniqueness of the solution of $(M_{i,n})$ that $\bar{u}_{i,n} = u_{i,n}$. \square

Along the same lines as in the proof of Theorem 2.2 in [19], we obtain the following uniform estimates:

Proposition 3.1. *Let $u_{i,n}^\varepsilon$ be the solution of $(M_{i,n}^\varepsilon)$. Then, for any $n \in \mathbb{N}$,*

$$(3.7) \quad \int_0^T \int_{\Omega} V_n^\varepsilon(x, t)^2 dx dt \leq 2E(u_0),$$

$$(3.8) \quad \sup_i \|\Delta u_{i,n}^\varepsilon\|_{L^2(\Omega)}^2 \leq 2E(u_0).$$

Since $N \leq 3$, combining Proposition 3.1 with Sobolev's embedding theorem, we have

$$(3.9) \quad u_{i,n}^\varepsilon \text{ is uniformly continuous in } \Omega, \text{ with modulus of continuity independent of } \varepsilon, i, \text{ and } n.$$

Set

$$\begin{aligned} \mathcal{C}_{i,n}^{\varepsilon,+} &= \{x \in \Omega \mid u_{i,n}^\varepsilon(x) = f_\varepsilon(x)\}, \\ \mathcal{C}_{i,n}^{\varepsilon,-} &= \{x \in \Omega \mid u_{i,n}^\varepsilon(x) = g(x)\}. \end{aligned}$$

By the fact that $f_\varepsilon < g$ in Ω , we observe from (3.9) that the sets $\mathcal{C}_{i,n}^{\varepsilon,+}$ and $\mathcal{C}_{i,n}^{\varepsilon,-}$ are disjoint. Here we set

$$\mu_{i,n}^\varepsilon = \Delta^2 u_{i,n}^\varepsilon + V_{i,n}^\varepsilon.$$

In the following, we show that $\mu_{i,n}^\varepsilon$ is a signed measure in Ω . To this aim, let us define

$$\gamma_\rho(\lambda) := \begin{cases} \frac{\lambda^2}{\rho} & \text{if } \lambda < 0, \\ 0 & \text{if } \lambda > 0, \end{cases}$$

$$\beta_\rho(\lambda) := \gamma'_\rho(\lambda),$$

for each $\rho > 0$. Regarding the following minimization problem

$$(M_{i,n}^{\varepsilon,\rho}) \quad \min_{v \in H_0^2(\Omega)} G_{i,n}^{\varepsilon,\rho}(v)$$

with

$$G_{i,n}^{\varepsilon,\rho}(v) := \int_{\Omega} \left[\frac{1}{2}(\Delta v)^2 + \frac{1}{2\tau_n}(v - u_{i-1,n}^\varepsilon)^2 + \gamma_\rho(v - f_\varepsilon) + \gamma_\rho(g - v) \right] dx,$$

we show the following:

Proposition 3.2. *The problem $(M_{i,n}^{\varepsilon,\rho})$ has a unique solution $w_{i,n}^{\varepsilon,\rho}$ with*

$$(3.10) \quad w_{i,n}^{\varepsilon,\rho} \rightharpoonup u_{i,n}^\varepsilon \quad \text{weakly in } H^2(\Omega) \quad \text{as } \rho \downarrow 0.$$

Proof. By a standard argument, we deduce that the problem $(M_{i,n}^{\varepsilon,\rho})$ has a unique solution $w_{i,n}^{\varepsilon,\rho}$ satisfying

$$\Delta^2 w_{i,n}^{\varepsilon,\rho} + \frac{1}{\tau_n}(w_{i,n}^{\varepsilon,\rho} - u_{i-1,n}^\varepsilon) + \beta_\rho(w_{i,n}^{\varepsilon,\rho} - f_\varepsilon) - \beta_\rho(g - w_{i,n}^{\varepsilon,\rho}) = 0 \quad \text{in } \Omega$$

in the classical sense. Since it follows from the minimality of $w_{i,n}^{\varepsilon,\rho}$ that

$$(3.11) \quad G_{i,n}^{\varepsilon,\rho}(w_{i,n}^{\varepsilon,\rho}) \leq G_{i,n}^{\varepsilon,\rho}(u_{i-1,n}^\varepsilon) = E(u_{i-1,n}^\varepsilon),$$

we observe from Proposition 3.1 that

$$(3.12) \quad \|\Delta w_{i,n}^{\varepsilon,\rho}\|_{L^2(\Omega)}^2 \leq 2E(u_0),$$

$$(3.13) \quad \frac{1}{2\tau_n} \|w_{i,n}^{\varepsilon,\rho} - u_{i-1,n}^\varepsilon\|_{L^2(\Omega)}^2 \leq E(u_0),$$

and

$$(3.14) \quad \max\{ \|(w_{i,n}^{\varepsilon,\rho} - f_\varepsilon)^-\|_{L^2(\Omega)}^2, \|(g - w_{i,n}^{\varepsilon,\rho})^-\|_{L^2(\Omega)}^2 \} \leq \rho E(u_0).$$

The inequality (3.12) yields that there exist a sequence $\{\rho_m\}$ with $\rho_m \rightarrow 0$ as $m \rightarrow \infty$ and a function $\tilde{u} \in H_0^2(\Omega)$ such that

$$(3.15) \quad w_{i,n}^{\varepsilon,\rho_m} \rightharpoonup \tilde{u} \quad \text{weakly in } H^2(\Omega),$$

in particular,

$$(3.16) \quad w_{i,n}^{\varepsilon,\rho_m} \rightarrow \tilde{u} \quad \text{a.e. in } \Omega,$$

as $\rho_m \rightarrow 0$. Recalling (3.14) and (3.16), we deduce from Chebyshev's inequality that $f_\varepsilon \leq \tilde{u} \leq g$ in Ω . This implies $\tilde{u} \in K_\varepsilon$.

We claim that \tilde{u} is a minimizer of $(M_{i,n}^\varepsilon)$. Indeed, for any $v \in K_\varepsilon$, it holds that

$$(3.17) \quad G_{i,n}^\varepsilon(v) = G_{i,n}^{\varepsilon,\rho_m}(v) \geq G_{i,n}^{\varepsilon,\rho_m}(w_{i,n}^{\varepsilon,\rho_m}) \geq G_{i,n}^\varepsilon(w_{i,n}^{\varepsilon,\rho_m}).$$

Recalling (3.15)-(3.16) and letting $\rho_m \rightarrow 0$ in (3.17), we infer that

$$G_{i,n}^\varepsilon(v) \geq \liminf_{\rho_m \downarrow 0} G_{i,n}^\varepsilon(w_{i,n}^{\varepsilon,\rho_m}) = G_{i,n}^\varepsilon(\tilde{u}).$$

This implies that \tilde{u} is a minimizer of $(M_{i,n}^\varepsilon)$. Then it follows from the uniqueness of the solutions to $(M_{i,n}^\varepsilon)$ that $\tilde{u} = u_{i,n}^\varepsilon$. We thus completed the proof. \square

Theorem 3.2. *Let $\varepsilon > 0$ and $i \in \{1, 2, \dots, n\}$. Then the quantity $\mu_{i,n}^\varepsilon$ is a signed measure in Ω with*

$$(3.18) \quad \text{supp } \mu_{i,n}^\varepsilon \subset \mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}, \quad \mu_{i,n}^\varepsilon \begin{cases} \geq 0 & \text{in } \mathcal{C}_{i,n}^{\varepsilon,+}, \\ \leq 0 & \text{in } \mathcal{C}_{i,n}^{\varepsilon,-}. \end{cases}$$

Moreover there exists a positive constant $C > 0$ independent of ε and n such that

$$(3.19) \quad \tau_n \sum_{i=1}^n \mu_{i,n}^\varepsilon(\Omega)^2 < C.$$

Proof. To begin with, we shall verify that the quantity

$$\mu_{i,n}^{\varepsilon,\rho} := \Delta^2 w_{i,n}^{\varepsilon,\rho} + (w_{i,n}^{\varepsilon,\rho} - u_{i-1,n}^\varepsilon)/\tau_n$$

defines a signed measure in Ω . Let us set

$$I_\rho^+ = \{x \in \Omega \mid w_{i,n}^{\varepsilon,\rho}(x) \leq f_\varepsilon(x)\}, \quad I_\rho^- = \{x \in \Omega \mid w_{i,n}^{\varepsilon,\rho}(x) \geq g(x)\}.$$

It follows from $\beta_\rho \leq 0$ that

$$\Delta^2 w_{i,n}^{\varepsilon,\rho} + \frac{w_{i,n}^{\varepsilon,\rho} - u_{i-1,n}^\varepsilon}{\tau_n} = -\beta_\rho(w_{i,n}^{\varepsilon,\rho} - f_\varepsilon) + \beta_\rho(g - w_{i,n}^{\varepsilon,\rho}) \begin{cases} \geq 0 & \text{in } I_\rho^+, \\ = 0 & \text{in } \Omega \setminus (I_\rho^+ \cup I_\rho^-), \\ \leq 0 & \text{in } I_\rho^-, \end{cases}$$

i.e., $\mu_{i,n}^{\varepsilon,\rho}$ defines a signed measure in Ω .

We claim that the measure $\mu_{i,n}^{\varepsilon,\rho}$ converges to $\mu_{i,n}^\varepsilon$ as $\rho \downarrow 0$ up to a subsequence. Indeed, we shall show that, for each ε , i , and n , the quantity $\mu_{i,n}^{\varepsilon,\rho}(U)$ is uniformly bounded with respect to ρ for any $U \subset\subset \Omega$. From now on, we write $\mu_{i,n}^{\varepsilon,\rho} = \nu_+^\rho - \nu_-^\rho$, where ν_\pm^ρ are positive measures with their support in I_ρ^\pm , respectively. For any $\varphi \in C_c^\infty(\Omega)$ with $\varphi \equiv 1$ in U and $0 \leq \varphi \leq 1$ elsewhere, we observe that

$$\begin{aligned} \nu_\pm^\rho(U) &\leq \int_U \varphi d\nu_\pm^\rho = \pm \int_U \left[\Delta w_{i,n}^{\varepsilon,\rho} \Delta \varphi + \frac{1}{\tau_n} (w_{i,n}^{\varepsilon,\rho} - u_{i-1,n}^\varepsilon) \varphi \right] dx \\ &\leq E(w_{i,n}^{\varepsilon,\rho})^{\frac{1}{2}} E(\varphi)^{\frac{1}{2}} + \sqrt{\frac{2}{\tau_n}} \left(\frac{1}{2\tau_n} \int_\Omega (w_{i,n}^{\varepsilon,\rho} - u_{i-1,n}^\varepsilon)^2 dx \right)^{\frac{1}{2}} \left(\int_\Omega \varphi^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

Since it follows from (3.11) that

$$(3.20) \quad \frac{1}{2\tau_n} \int_\Omega (w_{i,n}^{\varepsilon,\rho} - u_{i-1,n}^\varepsilon)^2 dx \leq E(u_{i-1,n}^\varepsilon) - E(w_{i,n}^{\varepsilon,\rho}),$$

we observe from (3.12) and (3.20) that

$$(3.21) \quad \nu_\pm^\rho(U) \leq C(U) \left[E(u_0)^{\frac{1}{2}} + \left(\frac{E(u_{i-1,n}^\varepsilon) - E(w_{i,n}^{\varepsilon,\rho})}{\tau_n} \right)^{\frac{1}{2}} \right].$$

Thus there exist a sequence $\{\rho_{m'}\} \subset \{\rho_m\}$ and measures $\bar{\mu}_\pm$ such that

$$(3.22) \quad \nu_\pm^{\rho_{m'}} \rightharpoonup \bar{\mu}_\pm \quad \text{as } m' \rightarrow \infty,$$

i.e., for any $\zeta \in C_c(\Omega)$

$$\int_{\Omega} \zeta d\nu_{\pm}^{\rho_{m'}} \rightarrow \int_{\Omega} \zeta d\bar{\mu}_{\pm} \quad \text{as } m' \rightarrow \infty,$$

where $\{\rho_m\}$ is the sequence obtained in the proof of Proposition 3.2. Since Proposition 3.2 asserts that

$$\begin{aligned} \int_{\Omega} \zeta d\nu_{\pm}^{\rho_{m'}} &= \pm \int_{\Omega} \left[\Delta w_{i,n}^{\varepsilon, \rho_{m'}} \Delta \zeta + \frac{1}{\tau_n} (w_{i,n}^{\varepsilon, \rho_{m'}} - u_{i-1,n}^{\varepsilon}) \zeta \right] dx \\ &\rightarrow \pm \int_{\Omega} [\Delta u_{i,n}^{\varepsilon} \Delta \zeta + V_{i,n}^{\varepsilon} \zeta] dx \quad \text{for any } \zeta \in C_c^2(\Omega) \quad \text{as } m' \rightarrow \infty, \end{aligned}$$

the relation (3.22) implies $\bar{\mu}_{\pm} = \pm(\Delta^2 u_{i,n}^{\varepsilon} + V_{i,n}^{\varepsilon})$, respectively. We claim that

$$(3.23) \quad \text{supp } \bar{\mu}_+ \subset \mathcal{C}_{i,n}^{\varepsilon,+}, \quad \text{supp } \bar{\mu}_- \subset \mathcal{C}_{i,n}^{\varepsilon,-}.$$

It is sufficient to show the former relation. Let $x_0 \in \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$ be chosen arbitrarily. Then there exist a neighborhood W of x_0 and a constant $\delta > 0$ such that

$$u_{i,n}^{\varepsilon}(x) - f_{\varepsilon}(x) > \delta \quad \text{in } W \subset \Omega.$$

Since $w_{i,n}^{\varepsilon, \rho_{m'}}$ uniformly converges to $u_{i,n}^{\varepsilon}$ as $m' \rightarrow \infty$, there exists a constant $M > 0$ such that for any $m' > M$

$$|w_{i,n}^{\varepsilon, \rho_{m'}} - u_{i,n}^{\varepsilon}| \leq \frac{\delta}{2} \quad \text{in } W.$$

Thus we deduce that, for any $m' > M$,

$$w_{i,n}^{\varepsilon, \rho_{m'}}(x) - f_{\varepsilon}(x) \geq (u_{i,n}^{\varepsilon} - f_{\varepsilon}(x)) - |w_{i,n}^{\varepsilon, \rho_{m'}} - u_{i,n}^{\varepsilon}| > \frac{\delta}{2},$$

i.e., $W \subset \Omega \setminus I_{\rho_{m'}}^+$ for any $m' > M$. Hence we see that for any $\zeta \in C_c^2(W)$

$$\int_{\Omega} \zeta d\bar{\mu}_+ = \lim_{m' \rightarrow \infty} \int_{\Omega} \zeta d\nu_+^{\rho_{m'}} = 0.$$

This is equivalent to the former relation in (3.23). Recalling that $\mathcal{C}_{i,n}^{\varepsilon,+}$ and $\mathcal{C}_{i,n}^{\varepsilon,-}$ are disjoint set, we observe that $\mu_{i,n}^{\varepsilon}$ is a signed measure satisfying (3.18).

We turn to the proof of (3.19). For any $U \subset\subset \Omega$, it follows from (3.21) that

$$\begin{aligned} \mu_{i,n}^{\varepsilon} &\leq \mu_{i,n}^{\varepsilon} \llcorner_{\mathcal{C}_{i,n}^{\varepsilon,+}} \leq C(U)E(u_0)^{\frac{1}{2}} + C(U) \liminf_{\rho_{m'} \downarrow 0} \left(\frac{E(u_{i-1,n}^{\varepsilon}) - E(w_{i,n}^{\varepsilon, \rho_{m'}})}{\tau_n} \right)^{\frac{1}{2}} \\ &= C(U)E(u_0)^{\frac{1}{2}} + C(U) \left(\frac{E(u_{i-1,n}^{\varepsilon}) - E(u_{i,n}^{\varepsilon})}{\tau_n} \right)^{\frac{1}{2}} \end{aligned}$$

and

$$\begin{aligned} \mu_{i,n}^{\varepsilon} &\geq \mu_{i,n}^{\varepsilon} \llcorner_{\mathcal{C}_{i,n}^{\varepsilon,-}} \geq -C(U)E(u_0)^{\frac{1}{2}} - C(U) \liminf_{\rho_{m'} \downarrow 0} \left(\frac{E(u_{i-1,n}^{\varepsilon}) - E(w_{i,n}^{\varepsilon, \rho_{m'}})}{\tau_n} \right)^{\frac{1}{2}} \\ &= -C(U)E(u_0)^{\frac{1}{2}} - C(U) \left(\frac{E(u_{i-1,n}^{\varepsilon}) - E(u_{i,n}^{\varepsilon})}{\tau_n} \right)^{\frac{1}{2}}. \end{aligned}$$

Multiplying τ_n and summing over $i = 0, 1, \dots, n$, we find

$$(3.24) \quad \tau_n \sum_{i=0}^n \mu_{i,n}^\varepsilon (U)^2 \leq C'(U)E(u_0)T + C'(U)(E(u_0) - E(u_{n,n})) \leq C'(U)(T+1)E(u_0).$$

It follows from the condition (1.2) that there exists a constant $\delta_* > 0$ such that

$$d(\partial\Omega, \mathcal{C}_{i,n}^{\varepsilon,\pm}) \geq \delta_*.$$

Thus it follows from (3.18) that $\text{supp } \mu_{i,n}^\varepsilon \subset \Omega_{\delta_*/2}$, where $\Omega_\rho := \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > \rho\}$. Letting $U = \Omega_{\delta_*/2}$, we obtain the conclusion. \square

We shall now prove the $C^{1,1}$ regularity of $u_{i,n}^\varepsilon$ in Ω . In the following, for each $h \in L^2(\Omega)$, we denote by $\Delta^{-1}h$ the solution of

$$\begin{cases} -\Delta w = h & \text{in } \Omega, \\ w = 0 & \text{on } \partial\Omega. \end{cases}$$

We start with the following lemma:

Lemma 3.2. *For each $\varepsilon > 0$, $n \in \mathbb{N}$, and $i \in \{1, \dots, n\}$, there exists a function $v_{i,n}^\varepsilon$ satisfying the following:*

- (a) $v_{i,n}^\varepsilon = \Delta u_{i,n}^\varepsilon + \Delta^{-1}V_{i,n}^\varepsilon$ a.e. in Ω ;
- (b) $v_{i,n}^\varepsilon$ is upper semicontinuous in $\Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,-}$. On the other hand, $v_{i,n}^\varepsilon$ is lower semicontinuous in $\Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$;
- (c) for any $x_0 \in \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,-}$ and any sequence of balls $B_\rho(x_0) \subset \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,-}$, it holds that

$$\frac{1}{|B_\rho(x_0)|} \int_{B_\rho(x_0)} v_{i,n}^\varepsilon dx \downarrow v_{i,n}^\varepsilon(x_0) \quad \text{as } \rho \downarrow 0.$$

On the other hand, for any $x_1 \in \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$ and any sequence of balls $B_\rho(x_1) \subset \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$, we have

$$\frac{1}{|B_\rho(x_1)|} \int_{B_\rho(x_1)} v_{i,n}^\varepsilon dx \uparrow v_{i,n}^\varepsilon(x_1) \quad \text{as } \rho \downarrow 0.$$

Proof. Let us set

$$v_{i,n}^{\varepsilon,\rho}(x) = \frac{1}{|B_\rho(x)|} \int_{B_\rho(x)} [\Delta u_{i,n}^\varepsilon(y) + \Delta^{-1}V_{i,n}^\varepsilon(y)] dy.$$

If $u_{i,n}^\varepsilon \in C^\infty(\Omega)$, then Green's formula yields that for each $x_0 \in \Omega$

$$(3.25) \quad \begin{aligned} \Delta u_{i,n}^\varepsilon(x_0) + \Delta^{-1}V_{i,n}^\varepsilon(x_0) &= \frac{1}{|\partial B_\rho(x_0)|} \int_{\partial B_\rho(x_0)} [\Delta u_{i,n}^\varepsilon + \Delta^{-1}V_{i,n}^\varepsilon] dS \\ &\quad - \int_{B_\rho(x_0)} [\Delta^2 u_{i,n}^\varepsilon(x) + V_{i,n}^\varepsilon(x)] G_\rho(x - x_0) dx, \end{aligned}$$

where G_ρ is Green's function defined by

$$(3.26) \quad G_\rho(r) = \begin{cases} \frac{1}{2}(r - \rho) & \text{if } N = 1, \\ \frac{1}{2\pi} \log \frac{\rho}{r} & \text{if } N = 2, \\ \frac{1}{N(N-2)\omega(N)}(r^{N-2} - \rho^{N-2}) & \text{if } N \geq 3. \end{cases}$$

We note that $\omega(N)$ denotes the volume of unit ball in \mathbb{R}^N . Thanks to (3.18) and the fact that $G_{\rho'} > G_\rho$ if $\rho' > \rho$, we observe from (3.25) that

$$(3.27) \quad v_{i,n}^{\varepsilon,\rho}(x_0) \leq v_{i,n}^{\varepsilon,\rho'}(x_0) \quad \text{if } \rho < \rho' \quad \text{and} \quad B_{\rho'}(x_0) \subset \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,-}$$

and

$$(3.28) \quad v_{i,n}^{\varepsilon,\rho}(x_0) \geq v_{i,n}^{\varepsilon,\rho'}(x_0) \quad \text{if } \rho < \rho' \quad \text{and} \quad B_{\rho'}(x_0) \subset \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$$

For general $u_{i,n}^\varepsilon \in H_0^2(\Omega)$, making use of the mollification of $\Delta u_{i,n}^\varepsilon + \Delta^{-1}V_{i,n}^\varepsilon$, we are able to verify (3.27) and (3.28). Hence it follows from (3.27) and (3.28) that

$$v_{i,n}^{\varepsilon,\rho}(x) \downarrow \bar{v}_{i,n}^\varepsilon(x) \quad \text{as } \rho \downarrow 0 \quad \text{in } \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,-}$$

and

$$v_{i,n}^{\varepsilon,\rho}(x) \uparrow \tilde{v}_{i,n}^\varepsilon(x) \quad \text{as } \rho \downarrow 0 \quad \text{in } \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+},$$

for some functions $\bar{v}_{i,n}^\varepsilon$ and $\tilde{v}_{i,n}^\varepsilon$.

Since $v_{i,n}^{\varepsilon,\rho}$ is continuous in Ω , setting

$$v_{i,n}^\varepsilon(x) = \begin{cases} \bar{v}_{i,n}^\varepsilon(x) & \text{if } x \in \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,-}, \\ \tilde{v}_{i,n}^\varepsilon(x) & \text{if } x \in \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}, \end{cases}$$

we deduce that $v_{i,n}^\varepsilon$ is upper semicontinuous in $\Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,-}$, and is lower semicontinuous in $\Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$. Recalling that $\Delta u_{i,n}^\varepsilon + \Delta^{-1}V_{i,n}^\varepsilon \in L^2(\Omega)$, we see that

$$v_{i,n}^{\varepsilon,\rho} \rightarrow \Delta u_{i,n}^\varepsilon + \Delta^{-1}V_{i,n}^\varepsilon \quad \text{as } \rho \downarrow 0 \quad \text{a.e. in } \Omega.$$

Therefore we conclude that $v_{i,n}^\varepsilon = \Delta u_{i,n}^\varepsilon + \Delta^{-1}V_{i,n}^\varepsilon$ a.e. in Ω . \square

Lemma 3.3. *For any $x_0 \in \mathcal{C}_{i,n}^{\varepsilon,+}$, it holds that*

$$(3.29) \quad v_{i,n}^\varepsilon(x_0) - \Delta^{-1}V_{i,n}^\varepsilon(x_0) \geq \Delta f(x_0).$$

On the other hand, for any $x_1 \in \mathcal{C}_{i,n}^{\varepsilon,-}$, we have

$$(3.30) \quad v_{i,n}^\varepsilon(x_1) - \Delta^{-1}V_{i,n}^\varepsilon(x_1) \leq \Delta g(x_1).$$

Proof. Since the proof of (3.29) is similar to the proof of Lemma 3.3 in [19], we shall prove the latter assertion. Let $x_1 \in \mathcal{C}_{i,n}^{\varepsilon,-}$. Since $\mathcal{C}_{i,n}^{\varepsilon,+}$ and $\mathcal{C}_{i,n}^{\varepsilon,-}$ are disjoint, it holds that $\mathcal{C}_{i,n}^{\varepsilon,-} \subset \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$. Then there exists a sequence $\{y_m\} \subset \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$ with $y_m \rightarrow x_1$ as $m \rightarrow \infty$ such that

$$(3.31) \quad u_{i,n}^\varepsilon(y_m) - g(y_m) \uparrow 0.$$

For each y_m , let ρ be small enough such that $B_{\rho,m} := \{y \in \mathbb{R}^N \mid |y - y_m| < \rho\} \subset \Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$. It follows from Green's formula that

$$(3.32) \quad u_{i,n}^\varepsilon(y_m) = \frac{1}{|\partial B_{\rho,m}|} \int_{\partial B_{\rho,m}} u_{i,n}^\varepsilon dS - \int_{B_{\rho,m}} \Delta u_{i,n}^\varepsilon(y) G_\rho(y_m - y) dy$$

and

$$(3.33) \quad g(y_m) = \frac{1}{|\partial B_{\rho,m}|} \int_{\partial B_{\rho,m}} g dS - \int_{B_{\rho,m}} \Delta g(y) G_\rho(y_m - y) dy,$$

Since $u_{i,n}^\varepsilon \leq g$ in Ω , we infer from (3.31)–(3.33) that

$$\liminf_{m \rightarrow \infty} \int_{B_{\rho,m}} [\Delta g(y) - \Delta u_{i,n}^\varepsilon(y)] G_\rho(y_m - y) dy \geq 0.$$

Thanks to Lemma 3.2, the relation is reduced to

$$(3.34) \quad \liminf_{m \rightarrow \infty} \int_{B_{\rho,m}} [\Delta g(y) - v_{i,n}^\varepsilon(y) + \Delta^{-1} V_{i,n}^\varepsilon(y)] G_\rho(y_m - y) dy \geq 0.$$

Recalling that $V_{i,n}^\varepsilon \in H_0^2(\Omega)$, we observe from the elliptic regularity, e.g., see [17], that $\Delta^{-1} V_{i,n}^\varepsilon \in H^4(\Omega)$. We note that Sobolev's embedding theorem implies that $\Delta^{-1} V_{i,n}^\varepsilon$ is continuous in Ω provided $N \leq 7$. Since $v_{i,n}^\varepsilon$ is lower semicontinuous in $\Omega \setminus \mathcal{C}_{i,n}^{\varepsilon,+}$, there exists a point $y_{m,\rho} \in \overline{B_{\rho,m}}$ such that the maximum of $\Delta g(y) - v_{i,n}^\varepsilon(y) + \Delta^{-1} V_{i,n}^\varepsilon(y)$ in $\overline{B_{\rho,m}}$ attains at $y = y_{m,\rho}$. Hence it follows from (3.34) that there exists a sequence $\{\delta_m\}$ with $\delta_m \downarrow 0$ as $m \rightarrow \infty$ such that

$$\Delta g(y_{m,\rho}) - v_{i,n}^\varepsilon(y_{m,\rho}) + \Delta^{-1} V_{i,n}^\varepsilon(y_{m,\rho}) \geq -\delta_m.$$

As $m \rightarrow \infty$, $y_{m,\rho}$ converges to a point $y_\rho \in \{y \in \mathbb{R}^N \mid |y - x_1| \leq \rho\}$ up to a subsequence, for the sequence $\{y_{m,\rho}\}$ is bounded. Thanks to the lower semicontinuity of $v_{i,n}^\varepsilon$, we find

$$\Delta g(y_\rho) - v_{i,n}^\varepsilon(y_\rho) + \Delta^{-1} V_{i,n}^\varepsilon(y_\rho) \geq 0$$

for any $\rho > 0$ small enough. Letting $\rho \downarrow 0$ and making use of the lower semicontinuity of $v_{i,n}^\varepsilon$, we conclude (3.30). \square

Lemma 3.4. *For each $\varepsilon > 0$, $n \in \mathbb{N}$, and $i = 1, \dots, n$, it holds that $\Delta u_{i,n}^\varepsilon \in L^\infty(\Omega)$. Moreover, there exists a positive constant C independent of ε , n , and i , such that*

$$(3.35) \quad \begin{aligned} \|\Delta u_{i,n}^\varepsilon\|_{L^\infty(\Omega)} &\leq CE(u_0)^{\frac{1}{2}} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \max\{\|\Delta f_\varepsilon\|_{L^\infty(\Omega)}, \|\Delta g\|_{L^\infty(\Omega)}\} \\ &\quad + C \left(\frac{E(u_{i-1,n}^\varepsilon) - E(u_{i,n}^\varepsilon)}{\tau_n} \right)^{\frac{1}{2}}. \end{aligned}$$

Proof. Let us set

$$(3.36) \quad U_{i,n}^\varepsilon := u_{i,n}^\varepsilon + (\Delta^2)^{-1} V_{i,n}^\varepsilon,$$

where $(\Delta^2)^{-1} V_{i,n}^\varepsilon$ denotes the unique solution of

$$\begin{cases} \Delta^2 w = V_{i,n}^\varepsilon & \text{in } \Omega, \\ w = 0, \Delta w = 0, & \text{on } \partial\Omega. \end{cases}$$

Fix $x_0 \in \Omega$ arbitrarily. Let B_ρ denote the ball center x_0 and the radius ρ . For any $R > 0$ with $\overline{B_R} \subset \Omega$, let $\zeta \in C_c^\infty(B_R)$ be a test function with $\zeta = 1$ in $B_{2R/3}$, $0 \leq \zeta \leq 1$

elsewhere. By the same argument as in the proof of Lemma 3.4 in [19], we see that for any $x \in B_{R/2}$

$$(3.37) \quad v_{i,n}^\varepsilon(x) = - \int_{B_{R/2}} G_R(x-y) d\mu_{i,n}^\varepsilon(y) - I_1(x) + \alpha(x)$$

with

$$I_1(x) := \int_{D_{R/2}} \zeta(y) G_R(x-y) \Delta^2 U_{i,n}^\varepsilon(y) dy.$$

and

$$|\alpha(x)| \leq C_1 \|\Delta U_{i,n}^\varepsilon\|_{L^2(\Omega)} \quad \text{in } B_{R/2}.$$

Here G_R is Green's function given by (3.26) with $\rho = R$. We note that for any $x \in B_{R/3}$

$$|I_1(x)| \leq C |\mu_{i,n}^\varepsilon|(D_{R/2}) \leq C_2 E(u_0)^{\frac{1}{2}} + C_3 \left(\frac{E(u_{i-1,n}^\varepsilon) - E(u_{i,n}^\varepsilon)}{\tau_n} \right)^{\frac{1}{2}},$$

where the constants C_2 and C_3 are independent of ε and n . Set

$$\tilde{G}_R(x) = \int_{B_{R/2}} G_R(x-y) d\mu_{i,n}^\varepsilon(y).$$

Thanks to Lemma 3.3, we observe from (3.37) that

$$\begin{aligned} \tilde{G}_R(x) &= -v_{i,n}^\varepsilon(x) - I_1(x) + \alpha(x) \leq -\Delta^{-1} V_{i,n}^\varepsilon(x) - \Delta f_\varepsilon(x) + |I_1(x)| + \alpha(x) \\ &< C_1 \|\Delta U_{i,n}^\varepsilon\|_{L^2(\Omega)} + C_2 E(u_0)^{\frac{1}{2}} + C_3 \left(\frac{E(u_{i-1,n}^\varepsilon) - E(u_{i,n}^\varepsilon)}{\tau_n} \right)^{\frac{1}{2}} \\ &\quad + C_4 \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|\Delta f_\varepsilon\|_{L^\infty(\Omega)} \quad \text{in } \mathcal{C}_{i,n}^{\varepsilon,+} \cap B_{R/3}, \end{aligned}$$

and while

$$\begin{aligned} \tilde{G}_R(x) &= -v_{i,n}^\varepsilon(x) - I_1(x) + \alpha(x) \geq -\Delta^{-1} V_{i,n}^\varepsilon(x) - \Delta g(x) - |I_1(x)| + \alpha(x) \\ &> -C_1 \|\Delta U_{i,n}^\varepsilon\|_{L^2(\Omega)} - C_2 E(u_0)^{\frac{1}{2}} - C_3 \left(\frac{E(u_{i-1,n}^\varepsilon) - E(u_{i,n}^\varepsilon)}{\tau_n} \right)^{\frac{1}{2}} \\ &\quad - C_4 \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} - \|\Delta g\|_{L^\infty(\Omega)} \quad \text{in } \mathcal{C}_{i,n}^{\varepsilon,-} \cap B_{R/3}. \end{aligned}$$

Then, along the same lines as in the proof of Theorems 1.6 and 1.10 of [18], we deduce that

$$\begin{aligned} \limsup_{d(x, \mathcal{C}_{i,n}^{\varepsilon,+}) \rightarrow 0} \tilde{G}_R(x) &\leq C_1 \|\Delta U_{i,n}^\varepsilon\|_{L^2(\Omega)} + C_2 E(u_0)^{\frac{1}{2}} + C_3 \left(\frac{E(u_{i-1,n}^\varepsilon) - E(u_{i,n}^\varepsilon)}{\tau_n} \right)^{\frac{1}{2}} \\ &\quad + C_4 \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|\Delta f_\varepsilon\|_{L^\infty(\Omega)} \end{aligned}$$

and

$$\begin{aligned} \limsup_{d(x, \mathcal{C}_{i,n}^{\varepsilon,-}) \rightarrow 0} \tilde{G}_R(x) &\geq -C_1 \|\Delta U_{i,n}^\varepsilon\|_{L^2(\Omega)} - C_2 E(u_0)^{\frac{1}{2}} - C_3 \left(\frac{E(u_{i-1,n}^\varepsilon) - E(u_{i,n}^\varepsilon)}{\tau_n} \right)^{\frac{1}{2}} \\ &\quad - C_4 \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} - \|\Delta g\|_{L^\infty(\Omega)}. \end{aligned}$$

Thus the maximal principle implies that

$$|\tilde{G}_R(x)| \leq C_1 \|\Delta U_{i,n}^\varepsilon\|_{L^2(\Omega)} + C_2 E(u_0)^{\frac{1}{2}} + C_3 \left(\frac{E(u_{i-1,n}^\varepsilon) - E(u_{i,n}^\varepsilon)}{\tau_n} \right)^{\frac{1}{2}} \\ + C_4 \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \max\{\|\Delta f_\varepsilon\|_{L^\infty(\Omega)}, \|\Delta g\|_{L^\infty(\Omega)}\} \quad \text{in } B_{R/3}.$$

Combining (3.37) with Theorem 3.2 and Lemma 3.2, we obtain (3.35). \square

Lemma 3.5. ([8]) *Let $N \leq 3$. Let $w \in H^2(\Omega)$ be a non-negative function satisfying*

$$\|\Delta w\|_{L^\infty(\Omega)} \leq M_0.$$

Then there exists a constant M depending only on M_0 such that if

$$x_0 \in J := \{x \in \Omega \mid w(x) = 0\}$$

then it holds that

$$(3.38) \quad |w(x)| \leq M|x - x_0|^2, \quad |\nabla w(x)| \leq M|x - x_0|, \quad \text{in } B(x_0, \rho/2),$$

where $\rho = \text{dist}(x_0, \partial\Omega)$.

Lemma 3.6. *For any $x \in \Omega \setminus (\mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-})$, it holds that*

$$|D^2 u_{i,n}^\varepsilon(x)| \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|D^2 f\|_{L^\infty(\Omega)} + \|\Delta^2 f\|_{L^2(\Omega)} \\ + \|D^2 g\|_{L^\infty(\Omega)} + \|\Delta^2 g\|_{L^2(\Omega)}).$$

Proof. Since $u_{i,n}^\varepsilon$ is continuous in Ω , we see that $\delta := \text{dist}(\mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}, \partial\Omega) > 0$. To begin with, recall that

$$\Delta^2 u_{i,n}^\varepsilon + V_{i,n}^\varepsilon = 0 \quad \text{in } \Omega \setminus \Omega_\delta,$$

where $\Omega_\delta = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > \delta\}$. By the elliptic regularity theory (i.e., see [17]), we deduce from $\partial\Omega \in C^4$ that

$$(3.39) \quad \|\Delta u_{i,n}^\varepsilon\|_{H^2(\Omega \setminus \Omega_\rho)} \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)}) \quad \text{for any } 0 < \rho < \delta,$$

where the constant $C > 0$ is independent of i , n and ε . Setting $\tilde{u} := \eta u_{i,n}^\varepsilon$, where $\eta \in C_c^\infty(\Omega \setminus \Omega_\delta)$ with $0 \leq \eta \leq 1$ and

$$\eta(x) = \begin{cases} 1 & \text{in } \Omega \setminus \Omega_{3\delta/4}, \\ 0 & \text{in } \Omega_{7\delta/8}, \end{cases}$$

we find

$$\begin{cases} \Delta^2 \tilde{u} = F(\eta, u_{i,n}^\varepsilon) - \eta V_{i,n}^\varepsilon & \text{in } \Omega \setminus \Omega_{7\delta/8}, \\ \tilde{u} = \partial_\nu \tilde{u} = 0 & \text{on } \partial(\Omega \setminus \Omega_{7\delta/8}), \end{cases}$$

where

$$F(\eta, u_{i,n}^\varepsilon) := \Delta^2 \eta u_{i,n}^\varepsilon + 2\nabla \Delta \eta \cdot \nabla u_{i,n}^\varepsilon + 2\Delta(\nabla \eta \cdot \nabla u_{i,n}^\varepsilon) + 2\Delta \eta \Delta u_{i,n}^\varepsilon + 2\nabla \eta \cdot \nabla \Delta u_{i,n}^\varepsilon.$$

Thanks to Theorem 2.20 in [14], we observe from (3.39) and $\partial\Omega \in C^4$ that

$$\|\tilde{u}\|_{H^4(\Omega \setminus \Omega_{7\delta/8})} \leq C(\|F(\eta, u_{i,n}^\varepsilon)\|_{L^2(\Omega \setminus \Omega_{7\delta/8})} + \|\eta V_{i,n}^\varepsilon\|_{L^2(\Omega \setminus \Omega_{7\delta/8})}) \\ \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)}).$$

Since $\|u_{i,n}^\varepsilon\|_{H^4(\Omega \setminus \Omega_{3\delta/4})} = \|\tilde{u}\|_{H^4(\Omega \setminus \Omega_{3\delta/4})} \leq \|\tilde{u}\|_{H^4(\Omega \setminus \Omega_{7\delta/8})}$, the estimate implies

$$\|D^2 u_{i,n}^\varepsilon\|_{H^2(\Omega \setminus \Omega_{3\delta/4})} \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)}).$$

Then it follows from Sobolev's embedding theorem that

$$(3.40) \quad \|D^2 u_{i,n}^\varepsilon\|_{L^\infty(\Omega \setminus \Omega_{3\delta/4})} \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)}),$$

where the constant C is independent of i , n , and ε .

Let $x_0 \in \Omega_{\delta/2} \setminus (\mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-})$ satisfy $\text{dist}(x_0, \mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}) \leq \delta$. Here we may assume that

$$\text{dist}(x_0, \mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}) = \text{dist}(x_0, \mathcal{C}_{i,n}^{\varepsilon,-}).$$

From Lemmas 3.4 and 3.5, there exists a constant $C > 0$ independent of i , n , and ε such that

$$(3.41) \quad |(u_{i,n}^\varepsilon - g)(x)| \leq C\|\Delta(u_{i,n}^\varepsilon - g)\|_{L^\infty(\Omega)}\text{dist}(x, \mathcal{C}_{i,n}^{\varepsilon,-})^2,$$

$$(3.42) \quad |\nabla(u_{i,n}^\varepsilon - g)(x)| \leq C\|\Delta(u_{i,n}^\varepsilon - g)\|_{L^\infty(\Omega)}\text{dist}(x, \mathcal{C}_{i,n}^{\varepsilon,-}),$$

in $B(x_0, d)$, where $d = \text{dist}(x_0, \mathcal{C}_{i,n}^{\varepsilon,-})$. We consider

$$w_d(x) = \frac{1}{d^2}(u_{i,n}^\varepsilon - g)(d(x - x_0)) \quad \text{in } B(x_0, 1).$$

For the simplicity, we may assume $x_0 = 0$. Then it follows from (3.41)-(3.42) that

$$|w_d(x)| \leq C\|\Delta(u_{i,n}^\varepsilon - g)\|_{L^\infty(\Omega)}, \quad |\nabla w_d(x)| \leq C\|\Delta(u_{i,n}^\varepsilon - g)\|_{L^\infty(\Omega)}, \quad \text{in } B(0, 1).$$

Since

$$\Delta^2 w_d(x) = -d^2 V_{i,n}^\varepsilon(d(x - x_0)) - d^2 \Delta^2 g(d(x - x_0)) \quad \text{in } B(0, 1),$$

we observe from the same argument as in the derivation of (3.40) that

$$|D^2 w_d(x)| \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \sum_{i=1}^2 \|\Delta^i g\|_{L^2(\Omega)}) \quad \text{in } B(0, \frac{1}{2}).$$

Thus it holds that

$$(3.43) \quad |D^2 u_{i,n}^\varepsilon(x)| \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|D^2 g\|_{L^\infty(\Omega)} + \|\Delta^2 g\|_{L^2(\Omega)}) \quad \text{in } B(x_0, d/2).$$

If $\text{dist}(x_0, \mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}) = \text{dist}(x_0, \mathcal{C}_{i,n}^{\varepsilon,+})$, then we obtain (3.43) replaced g by f . We thus completed the proof. \square

Theorem 3.3. *It holds that $u_{i,n}^\varepsilon \in W^{2,\infty}(\Omega)$. Moreover, there exists a positive constant C independent of ε and n such that*

$$\begin{aligned} \tau_n \sum_{i=1}^n \|D^2 u_{i,n}^\varepsilon\|_{L^\infty(\Omega)}^2 &\leq C(E(u_0) + \|D^2 f\|_{L^\infty(\Omega)} + \|\Delta^2 f\|_{L^2(\Omega)} \\ &\quad + \|D^2 g\|_{L^\infty(\Omega)} + \|\Delta^2 g\|_{L^2(\Omega)}). \end{aligned}$$

Proof. Let e_j be the unit vector in the direction of the positive x_j axis. Fix $x \in \Omega$. For $|h| \in \mathbb{R}$ small enough, we consider the second order differential quotient

$$D_h^2 u_{i,n}^\varepsilon(x) = \frac{u_{i,n}^\varepsilon(x + he_j) + u_{i,n}^\varepsilon(x - he_j) - 2u_{i,n}^\varepsilon(x)}{2h^2}.$$

If $\text{dist}(x, \mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}) < 4|h|$, then there exists $x_0 \in \mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}$ such that

$$|x - x_0| = \text{dist}(x, \mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}) < 4|h|.$$

We may assume $x_0 \in \mathcal{C}_{i,n}^{\varepsilon,-}$. Making use of (3.41), we find

$$\begin{aligned} & |D_h^2(u_{i,n}^\varepsilon - g)(x)| \\ & \leq \frac{C}{h^2} \|\Delta(u_{i,n}^\varepsilon - g)\|_{L^\infty(\Omega)} [\text{dist}(x + he_j, \mathcal{C}_{i,n}^{\varepsilon,-})^2 + \text{dist}(x - he_j, \mathcal{C}_{i,n}^{\varepsilon,-})^2 + \text{dist}(x, \mathcal{C}_{i,n}^{\varepsilon,-})^2] \\ & \leq C \|\Delta(u_{i,n}^\varepsilon - g)\|_{L^\infty(\Omega)}. \end{aligned}$$

On the other hand, if $\text{dist}(x, \mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}) \geq 4|h|$, then we observe from Lemma 3.6 that

$$\begin{aligned} |D_h^2 u_{i,n}^\varepsilon(x)| & \leq |D_{x_j x_j} u_{i,n}^\varepsilon(\tilde{x})| \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|D^2 f\|_{L^\infty(\Omega)} \\ & \quad + \|\Delta^2 f\|_{L^2(\Omega)} + \|D^2 g\|_{L^\infty(\Omega)} + \|\Delta^2 g\|_{L^2(\Omega)}), \end{aligned}$$

where $\tilde{x} \in B(x, 2\text{dist}(x, \mathcal{C}_{i,n}^{\varepsilon,+} \cup \mathcal{C}_{i,n}^{\varepsilon,-}))$. Consequently we see that, for any $x \in \Omega$, if $|h|$ is small enough,

$$\begin{aligned} |D_h^2 u_{i,n}^\varepsilon(x)| & \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^\infty(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|D^2 f\|_{L^\infty(\Omega)} \\ & \quad + \|\Delta^2 f\|_{L^2(\Omega)} + \|D^2 g\|_{L^\infty(\Omega)} + \|\Delta^2 g\|_{L^2(\Omega)}), \end{aligned}$$

where $C > 0$ is independent of x and h . Therefore we deduce that

$$(3.44) \quad \begin{aligned} |D_{x_j x_j} u_{i,n}^\varepsilon(x)| & \leq C(\|\Delta u_{i,n}^\varepsilon\|_{L^\infty(\Omega)} + \|V_{i,n}^\varepsilon\|_{L^2(\Omega)} + \|D^2 f\|_{L^\infty(\Omega)} \\ & \quad + \|\Delta^2 f\|_{L^2(\Omega)} + \|D^2 g\|_{L^\infty(\Omega)} + \|\Delta^2 g\|_{L^2(\Omega)}) \quad \text{in } \Omega. \end{aligned}$$

Combining (3.44) with Proposition 3.1 and Lemma 3.4, we obtain the conclusion. \square

Let us set

$$(3.45) \quad \mathcal{C}_{i,n}^+ = \{x \in \Omega \setminus \Omega_0 \mid u_{i,n}(x) = f(x)\},$$

$$(3.46) \quad \mathcal{C}_{i,n}^- = \{x \in \Omega \setminus \Omega_0 \mid u_{i,n}(x) = g(x)\},$$

where Ω_0 is defined in (1.6).

Theorem 3.4. *As $\varepsilon \downarrow 0$, the signed measure $\mu_{i,n}^\varepsilon$ converges to a signed Radon measure $\mu_{i,n}$ in Ω defined by*

$$\mu_{i,n} = \begin{cases} \Delta^2 u_{i,n} + V_{i,n} & \text{in } \Omega \setminus \Omega_0, \\ \Delta^2 f & \text{in } \Omega_0. \end{cases}$$

Moreover it holds that $\text{supp } \mu_{i,n} \subset \mathcal{C}_{i,n}^+ \cup \mathcal{C}_{i,n}^- \cup \Omega_0$,

$$\mu_{i,n} \begin{cases} \geq 0 & \text{in } \mathcal{C}_{i,n}^+, \\ \leq 0 & \text{in } \mathcal{C}_{i,n}^-, \end{cases}$$

and there exists a positive constant $C > 0$ independent of n such that

$$(3.47) \quad \tau_n \sum_{i=1}^n \mu_{i,n}(\Omega)^2 < CE(u_0) + T \|\Delta^2 f\|_{L^\infty(\Omega_0)}^2.$$

Proof. To begin with, we shall prove that $\mu_{i,n}^\varepsilon|_{\Omega_0} \rightharpoonup \Delta^2 f$ as $\varepsilon \downarrow 0$, i.e.,

$$(3.48) \quad \int_{\Omega} [\Delta u_{i,n}^\varepsilon \Delta \varphi + V_{i,n}^\varepsilon \varphi] dx \rightarrow \int_{\Omega} \Delta f \Delta \varphi dx \quad \text{as } \varepsilon \downarrow 0 \quad \text{for any } \varphi \in C_c^\infty(\Omega_0).$$

Since it holds that

$$|u_{i,n}^\varepsilon(x) - f(x)| \leq \varepsilon \quad \text{in } \Omega_0,$$

we infer that

$$(3.49) \quad \left| \int_{\Omega} (\Delta u_{i,n}^\varepsilon - \Delta f) \Delta \varphi dx \right| \leq \|u_{i,n}^\varepsilon - f\|_{L^\infty(\Omega_0)} \int_{\Omega} |\Delta^2 \varphi| dx \leq \varepsilon \int_{\Omega} |\Delta^2 \varphi| dx.$$

On the other hand, from

$$|V_{i,n}^\varepsilon| \leq \frac{1}{\tau_n} \{|u_{i,n}^\varepsilon - f| + |u_{i-1,n}^\varepsilon - f|\} \leq \frac{2}{\tau_n} \varepsilon,$$

we have

$$(3.50) \quad \left| \int_{\Omega} V_{i,n}^\varepsilon \varphi dx \right| \leq \frac{2}{\tau_n} \varepsilon \int_{\Omega} |\varphi| dx.$$

Then (3.49) and (3.50) implies (3.48).

From now on, we write $\mu_{i,n}^\varepsilon|_{\Omega \setminus \Omega_0} = \nu_{i,n}^{\varepsilon,+} - \nu_{i,n}^{\varepsilon,-}$, where $\nu_{i,n}^{\varepsilon,\pm}$ are positive measure in Ω with $\text{supp } \nu_{i,n}^{\varepsilon,\pm} \subset \mathcal{C}_{i,n}^{\varepsilon,\pm}$, respectively. By the proof of Theorem 3.2, there exist measures $\bar{\mu}_{i,n}^\pm$ in Ω such that

$$\nu_{i,n}^{\varepsilon,\pm} \rightharpoonup \bar{\mu}_{i,n}^\pm \quad \text{as } \varepsilon \downarrow 0,$$

i.e.,

$$\int_{\Omega} \zeta d\nu_{i,n}^{\varepsilon,\pm} \rightarrow \int_{\Omega} \zeta d\bar{\mu}_{i,n}^\pm \quad \text{for any } \zeta \in C_c(\Omega \setminus \Omega_0) \quad \text{as } \varepsilon \downarrow 0.$$

Since

$$\int_{\Omega} \zeta d\nu_{i,n}^{\varepsilon,\pm} = \pm \int_{\Omega} [\Delta u_{i,n}^\varepsilon \Delta \zeta + V_{i,n}^\varepsilon \zeta] dx \rightarrow \pm \int_{\Omega} [\Delta u_{i,n} \Delta \zeta + V_{i,n} \zeta] dx$$

for any $\zeta \in C_c^2(\Omega \setminus \Omega_0)$ as $\varepsilon \downarrow 0$, it holds that $\bar{\mu}_{i,n}^\pm = \pm(\Delta^2 u_{i,n} + V_{i,n})$. We claim that

$$(3.51) \quad \text{supp } \bar{\mu}_{i,n}^+ \subset \mathcal{C}_{i,n}^+, \quad \text{supp } \bar{\mu}_{i,n}^- \subset \mathcal{C}_{i,n}^-.$$

It is sufficient to show the former relation. Let $x_0 \in \Omega \setminus (\mathcal{C}_{i,n}^+ \cup \Omega_0)$. Then there exist a neighborhood $W \subset \Omega \setminus \Omega_0$ of x_0 and a constant $\delta > 0$ such that

$$u_{i,n}(x) - f(x) > \delta \quad \text{in } W.$$

Since $u_{i,n}^\varepsilon$ uniformly converges to $u_{i,n}$, there exists $\varepsilon_* > 0$ such that for any $\varepsilon < \varepsilon_*$

$$|u_{i,n}^\varepsilon(x) - u_{i,n}(x)| < \frac{\delta}{3} \quad \text{in } W.$$

Thus, for any $\varepsilon < \min\{\varepsilon_*, \delta/3\}$, we have

$$u_{i,n}^\varepsilon(x) - f_\varepsilon(x) > u_{i,n}(x) - f(x) - |u_{i,n}^\varepsilon(x) - u_{i,n}(x)| - |f_\varepsilon(x) - f(x)| > \frac{\delta}{3} \quad \text{in } W,$$

i.e., $W \subset \Omega \setminus (\mathcal{C}_{i,n}^{\varepsilon,+} \cup \Omega_0)$ for $\varepsilon > 0$ small enough. Hence we infer that for any $\zeta \in C_c(W)$

$$\int_{\Omega} \zeta d\bar{\mu}_{i,n}^+ = \lim_{\varepsilon \downarrow 0} \int_{\Omega} \zeta d\nu_{i,n}^{\varepsilon,+} = 0.$$

Therefore the relation (3.51) holds.

Finally we turn to (3.47). It follows from the proof of Theorem 3.2 that

$$\bar{\mu}_{i,n}^{\pm}(\Omega) \leq \liminf_{\varepsilon \downarrow 0} \nu_{i,n}^{\varepsilon,\pm}(\Omega) \leq C(U)E(u_0)^{\frac{1}{2}} + C(U) \left(\frac{E(u_{i-1,n}^{\varepsilon}) - E(u_{i,n}^{\varepsilon})}{\tau_n} \right)^{\frac{1}{2}}.$$

Moreover it holds that

$$\tau_n \sum_{i=1}^n \mu_{i,n}|_{\Omega_0}(x)^2 = \tau_n \sum_{i=1}^n |\Delta^2 f(x)|^2 = T |\Delta^2 f(x)|^2 \leq T \|\Delta^2 f\|_{L^\infty(\Omega_0)}^2 \quad \text{in } \Omega_0.$$

Recalling that $\text{supp } \mu_{i,n} \subset \mathcal{C}_{i,n}^+ \cup \mathcal{C}_{i,n}^- \cup \Omega_0$, we obtain

$$\begin{aligned} \tau_n \sum_{i=1}^n \mu_{i,n}(\Omega)^2 &\leq C_1 T E(u_0) + C_2 \sum_{i=1}^n \{E(u_{i-1,n}^{\varepsilon}) - E(u_{i,n}^{\varepsilon})\} + T \|\Delta^2 f\|_{L^\infty(\Omega_0)}^2 \\ &\leq C E(u_0) + T \|\Delta^2 f\|_{L^\infty(\Omega_0)}^2. \end{aligned}$$

We thus completed the proof. □

4. PROOF OF THE MAIN THEOREM

In this section, we prove Theorem 1.1. First we shall prove the convergence of the piecewise linear interpolation u_n of $\{u_{i,n}\}$. The proof is followed from the uniform estimates on $\{u_n\}$. Since the estimates have already obtained by Proposition 3.1, we are able to prove the following result along the same lines as in the proof of Theorem 4.1 in [19].

Theorem 4.1. *Let u_n be the piecewise linear interpolation of $\{u_{i,n}\}$. Then there exists a function*

$$u \in L^\infty(0, T; H_0^2(\Omega)) \cap H^1(0, T; L^2(\Omega))$$

such that for any $T < \infty$

$$u_n \rightharpoonup u \quad \text{in } L^2(0, T; H^2(\Omega)) \cap H^1(0, T; L^2(\Omega)) \quad \text{as } n \rightarrow \infty,$$

up to a subsequence. Moreover

$$\int_0^T \int_{\Omega} |\partial_t u|^2 \, dx dt \leq 2E(u_0),$$

$f(x) \leq u(x, t) \leq g(x)$ for $x \in \Omega$ and every $t \in [0, T]$, and for each $\alpha \in (0, 1/2)$, it holds that

$$u_n \rightarrow u \quad \text{in } C^{0,\alpha}([0, T]; L^2(\Omega)) \quad \text{as } n \rightarrow \infty.$$

Next we investigate the regularity of the limit u obtained by Theorem 4.1. The proof depends only on the uniform estimate on u_n obtained in Theorem 3.3. The same argument as in the proof of Theorems 4.2 and 4.3 in [19] gives us the following:

Theorem 4.2. *Let u be the function obtained by Theorem 4.1. Then it holds that*

$$u_n \rightarrow u \quad \text{weakly}^* \text{ in } L^2(0, T; W^{2, \infty}(\Omega)) \quad \text{as } n \rightarrow \infty.$$

Moreover, if $N = 1$,

$$u_n \rightarrow u \quad \text{in } C^{0, \beta}([0, T]; C^{1, \alpha}(\Omega)) \quad \text{as } n \rightarrow \infty$$

for every $\alpha \in (0, 1/2)$ and $\beta \in (0, (1 - 2\alpha)/8)$, and if $N = 2, 3$,

$$u_n \rightarrow u \quad \text{in } C^{0, \beta}([0, T]; C^{0, \alpha}(\Omega)) \quad \text{as } n \rightarrow \infty$$

for every

$$0 < \alpha < 2 - \frac{N}{2}, \quad 0 < \beta < \left(\frac{1}{2} - \frac{N}{8} \right) \left(1 - \frac{\alpha}{2 - N/2} \right).$$

In order to complete the proof of Theorem 1.1, we make use of the convergence result on the piecewise constant interpolation of $\{u_{i,n}\}$.

Lemma 4.1. ([19]) *Let \tilde{u}_n be the piecewise constant interpolation of $\{u_{i,n}\}$. If $N = 1$,*

$$\tilde{u}_n \rightarrow u \quad \text{in } L^\infty(0, T; C^{1, \alpha}(\Omega)) \quad \text{as } n \rightarrow \infty$$

for every $\alpha \in (0, 1/2)$, where u is the function obtained by Theorem 4.1. If $N = 2, 3$,

$$\tilde{u}_n \rightarrow u \quad \text{in } L^\infty(0, T; C^{0, \alpha}(\Omega)) \quad \text{as } n \rightarrow \infty$$

for every $\alpha \in (0, 2 - N/2)$. Moreover, for any $N \geq 1$, it holds that

$$\Delta \tilde{u}_n \rightharpoonup \Delta u \quad \text{in } L^2(0, T; L^2(\Omega)) \quad \text{as } n \rightarrow \infty.$$

We are in a position to complete the proof of Theorem 1.1. Let us define

$$(4.1) \quad \mu_n(t) = \mu_{i,n} \quad \text{if } t \in ((i-1)\tau_n, i\tau_n],$$

and set

$$(4.2) \quad \mathcal{C}_f = \{ (x, t) \in (\Omega \setminus \Omega_0) \times \mathbb{R}_+ \mid u(x, t) = f(x) \},$$

$$(4.3) \quad \mathcal{C}_g = \{ (x, t) \in (\Omega \setminus \Omega_0) \times \mathbb{R}_+ \mid u(x, t) = g(x) \}.$$

Proof of Theorem 1.1 Let u be the function obtained by Theorem 4.1. To begin with, along the same lines as in [19], we see that

$$(4.4) \quad \int_0^T \int_\Omega [\partial_t u(w - u) + \Delta u \Delta(w - u)] dx dt \geq 0 \quad \text{for any } w \in \mathcal{K},$$

i.e., u is a weak solution of (P). Moreover the uniqueness follows from the results in [4].

By virtue of Theorem 3.4, we deduce that

$$\int_0^T \mu_n(\Omega)^2 dt = \sum_{i=1}^n \int_{(i-1)\tau_n}^{i\tau_n} \mu_{i,n}(\Omega)^2 dt = \tau_n \sum_{i=1}^n \mu_{i,n}(\Omega)^2 < CE(u_0) + T \|\Delta^2 f\|_{L^\infty(\Omega_0)}^2.$$

Thus, as $n \rightarrow \infty$,

$$\mu_n \rightharpoonup \bar{\mu} \quad \text{weakly in } L^2(0, T; \mathcal{M}(\Omega)),$$

i.e.,

$$\int_0^T \int_\Omega \varphi d\mu_n dt \rightarrow \int_0^T \int_\Omega \varphi d\bar{\mu} dt \quad \text{for any } \varphi \in L^2(0, T; C_c^\infty(\Omega)) \quad \text{as } n \rightarrow \infty.$$

Since $\mu_{i,n}|_{\Omega_0} = \Delta^2 f$, we observe from the definition of μ_n that

$$\mu_n(t)|_{\Omega_0} = \Delta^2 f \quad \text{in } [0, T) \quad \text{for any } n \in \mathbb{N}.$$

From now on, we set $\mu_n|_{\Omega \setminus \Omega_0} = \nu_n^+ - \nu_n^-$ with

$$\nu_n^\pm(t) = \mu_{i,n}^\pm \quad \text{if } t \in ((i-1)\tau_n, i\tau_n],$$

where $\mu_{i,n}^+$ and $\mu_{i,n}^-$ denote respectively the positive part and the negative part of $\mu_{i,n}|_{\Omega \setminus \Omega_0}$. Since Theorem 3.4 deduces that

$$\int_0^T \nu_n^\pm(\Omega)^2 dt < C,$$

there exist measures $\bar{\mu}_\pm$ such that

$$\nu_n^\pm \rightharpoonup \bar{\mu}_\pm \quad \text{weakly in } L^2(0, T; \mathcal{M}(\Omega)) \quad \text{as } n \rightarrow \infty,$$

i.e., for any $\varphi \in L^2(0, T; C_c^\infty(\Omega \setminus \Omega_0))$,

$$\int_0^T \int_\Omega \varphi d\nu_n^\pm dt \rightarrow \int_0^T \int_\Omega \varphi d\bar{\mu}_\pm dt \quad \text{as } n \rightarrow \infty.$$

On the other hand, it holds that

$$\begin{aligned} \int_0^T \int_\Omega \varphi d\nu_n^\pm dt &= \pm \int_0^T \int_\Omega [\Delta \tilde{u}_n \Delta \varphi + V_n \varphi] dx dt \\ &\rightarrow \pm \int_0^T \int_\Omega [\Delta u \Delta \varphi + \partial_t u \varphi] dx dt \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Thus we infer that $\bar{\mu}_\pm = \pm(\Delta^2 u + \partial_t u)$. We claim that

$$(4.5) \quad \text{supp } \bar{\mu}_+ \subset \mathcal{C}_f, \quad \text{supp } \bar{\mu}_- \subset \mathcal{C}_g.$$

We shall prove the former relation. Let $x_0 \in \Omega \setminus (\mathcal{C}_f \cup \Omega_0)$. Since u is continuous in $\Omega \times \mathbb{R}_+$, there exist an open set $W \subset \Omega \setminus \Omega_0$, $0 < t_1 < t_2 < T$, and $\delta > 0$ such that

$$u(x, t) - f(x) > \delta \quad \text{in } W \times (t_1, t_2).$$

It follows from Lemma 4.1 that there exists a constant $N > 0$ such that

$$\tilde{u}_n(x, t) - u(x, t) > -\frac{\delta}{2} \quad \text{in } W \times (t_1, t_2) \quad \text{for any } n \geq N,$$

so that

$$\tilde{u}_n(x, t) - f(x) > \frac{\delta}{2} \quad \text{in } W \times (t_1, t_2) \quad \text{for any } n \geq N.$$

This means that, for any $n \geq N$,

$$(4.6) \quad W \times (t_1, t_2) \subset \Omega \setminus (\mathcal{C}_{i,n}^+ \cup \Omega_0) \quad \text{for each } \left[\frac{t_1}{\tau_n} \right] \leq i \leq \left[\frac{t_2}{\tau_n} \right].$$

Thus we deduce that for any $\varphi \in C_c((t_1, t_2); C_c^\infty(W))$

$$\begin{aligned} \int_0^T \int_\Omega \varphi d\bar{\mu}_+ dt &= \lim_{n \rightarrow \infty} \int_0^T \int_\Omega \varphi d\nu_n^+ dt = \lim_{n \rightarrow \infty} \int_0^T \int_\Omega [\Delta \tilde{u}_n \Delta \varphi + V_n \varphi] dx dt \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \int_{(i-1)\tau_n}^{i\tau_n} \int_\Omega [\Delta u_{i,n} \Delta \varphi + V_{i,n} \varphi] dx dt = 0. \end{aligned}$$

The last equality follows from (4.6). This implies the relation (4.5). □

REFERENCES

- [1] L. Ambrosio, *Minimizing movements*, Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. (5) **19** (1995), 191–246.
- [2] G. Barbatis, *Explicit estimates on the fundamental solution of higher-order parabolic equations with measurable coefficients*, J. Differential Equations **174** (2001), 442–463.
- [3] C. Baiocchi, F. Gastaldi and F. Tomarelli, *Some existence results on noncoercive variational inequalities*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (4) **13** (1986), no.4, 617–659.
- [4] H. Brezis, *Opérateurs Maximaux Monotones et Semi-groupes de Contractions dans les Espaces de Hilbert*, North-Holland, Amsterdam/London, 1973.
- [5] H. Brezis and G. Stampacchia, *Remarks on some fourth order variational inequalities*, Ann. Sc. Norm. Super. Pisa Cl. Sci. **2** (1977), 363–371.
- [6] L. A. Caffarelli, *The obstacle problem revisited*, Jour. Fourier Anal. Appl. **4** (1998), no. 4-5, 383–402.
- [7] L. A. Caffarelli and A. Friedman, *The obstacle problem for the biharmonic operator*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (4) **6** (1979), 151–184.
- [8] L. A. Caffarelli, A. Friedman and A. Torelli, *The two-obstacle problem for the biharmonic operator*, Pacific J. Math. **103** (1982), no. 2, 325–335.
- [9] G. Caristi and E. Mitidieri, *Existence and nonexistence of global solutions of higher-order parabolic problems with slow decay initial data*, J. Math. Anal. Appl. **279** (2003), 710–722.
- [10] J. W. Cholewa and A. Rodriguez-Bernal, *Linear and semilinear higher order parabolic equations in \mathbb{R}^N* , Nonlinear Anal. **75** (2012), 194–210.
- [11] J. Frehse, *On the regularity of the solution of the biharmonic variational inequality*, Manuscripta Math. **9** (1973), 91–103.
- [12] F. Gazzola, *On the moments of solutions to linear parabolic equations involving the biharmonic operator*, Discrete Contin. Dyn. Syst. **33** (2013), no. 8, 3583–3597.
- [13] F. Gazzola and H.-C. Grunau, *Eventual local positivity for a biharmonic heat equation in \mathbb{R}^n* , Discrete Contin. Dyn. Syst. Ser. S **1** (2008), no. 1, 83–87.
- [14] F. Gazzola, H.-C. Grunau, G. Sweers, *Polyharmonic boundary value problems. Positivity preserving and nonlinear higher order elliptic equations in bounded domains*, Lecture Notes in Mathematics **1991**, Springer-Verlag, Berlin, 2010.
- [15] V. A. Galaktionov and P. J. Harwin, *Non-uniqueness and global similarity solutions for a higher-order semilinear parabolic equation*, Nonlinearity **18** (2005), 717–746.
- [16] V. A. Galaktionov and S. I. Pohozaev, *Existence and blow-up for higher-order semilinear parabolic equations: majorizing order-preserving operators*, Indiana Univ. Math. J. **51** (2002), no. 6, 1321–1338.
- [17] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, 1998.
- [18] N. S. Landkof, *Foundations of Modern Potential Theory*, Springer-Verlag, New York, 1972.
- [19] M. Novaga and S. Okabe, *Regularity of the obstacle problem for the parabolic biharmonic equation*, Math. Ann., to appear.
- [20] B. Schild, *On the coincident set in biharmonic variational inequalities with thin obstacle*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (4) **13** (1986), no.4, 559–616.

DIPTAMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, PISA, ITALY
E-mail address: novaga@dm.unipi.it

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, 980-8578 SENDAI, JAPAN
E-mail address: okabe@math.tohoku.ac.jp