

HORN MAPS OF SEMI-PARABOLIC HÉNON MAPS

MATTHIEU ASTORG AND FABRIZIO BIANCHI

ABSTRACT. We prove that horn maps associated to quadratic semi-parabolic fixed points of Hénon maps, first introduced by Bedford, Smillie, and Ueda, satisfy a weak form of the Ahlfors island property. As a consequence, two natural definitions of their Julia set (the non-normality locus of the family of iterates and the closure of the set of the repelling periodic points) coincide. As another consequence, we also prove that there exist small perturbations of semi-parabolic Hénon maps for which the Hausdorff dimension of the forward Julia set J^+ is arbitrarily close to 4.

1. INTRODUCTION

Following the seminal work of Douady-Hubbard [DH84, DH85a] and Lavaurs [Lav89], the study of perturbations of maps with a parabolic fixed point, often referred to as *parabolic implosion*, has been a major theme in one-variable complex dynamics. A first consequence of this theory is the discontinuity of the Julia sets at parameters where a parabolic bifurcation occurs [Dou94]. Further notable applications include the celebrated result by Shishikura [Shi98] that parabolic maps can be approximated by hyperbolic maps with arbitrarily large hyperbolic dimension and, as a consequence, the proof that the boundary of the Mandelbrot set has maximal Hausdorff dimension, see also [DSZ97, McM00, Tan98, Zin98] for refinements of this result and further consequences and [Shi00, PV20] for an overview of the theory.

Shishikura's proof involves the so-called *horn maps*. These maps describe the limit behaviour of the return maps of large iterates of the perturbed maps near the parabolic point. Another important aspect of horn maps (also called *Écalle-Voronin invariants*) is that they classify parabolic germs up to analytic conjugacy, i.e., they form a *complete invariant* for this notion of equivalence [Éca85, Vor81]. Techniques from parabolic implosion have been extended by Inou and Shishikura [IS06], who introduced the *near-parabolic renormalization*, a powerful tool which was used, for instance, to construct quadratic Julia sets with positive area [BC12], and to make significant steps towards the settling of the Fatou hyperbolicity conjecture [CS15].

In recent years, techniques of parabolic implosion have started to be developed and successfully applied also in higher dimensions. In [BSU17], Bedford, Smillie, and Ueda extended Lavaurs' results to diffeomorphisms of $(\mathbb{C}^2, 0)$ with a semi-parabolic fixed point, that is, with one multiplier equal to 1 and one in the unit disk. In the important particular case of dissipative Hénon map, the authors introduced a horn map analogous to the one-dimensional case, which they used to show the discontinuity of various dynamically meaningful sets at parameters with a semi-parabolic fixed point. In [DL15], Dujardin and Lyubich adapted and improved the results from [BSU17] to construct homoclinic tangencies in some regions of the parameter space of complex Hénon maps, as part of

their characterization of stability and bifurcation in families of such maps. Parabolic implosion techniques in two complex variables were also used by Buff, Dujardin, Peter, Raissy, and the first author to give the first example of an endomorphism of $\mathbb{P}^2(\mathbb{C})$ with a wandering domain [ABD⁺16], which solved a long-standing open question in the domain, see also [ABTP23]. Adaptations of these techniques from Boc Thaler and the first author have lead to a precise description of the local dynamics near a parabolic point of a significant class of maps tangent to the identity in \mathbb{C}^2 [Aba01], in particular solving a long-standing open question by Abate [Aba05] on the topological classification of such maps. The first result on the parabolic implosion of a two-dimensional map tangent to the identity was also established by the second author in [Bia19].

Coming back to the original work by Bedford, Smillie, and Ueda, very little is known about the dynamics of the horn maps of semi-parabolic Hénon maps; for instance, it was not even known until now whether they always had periodic cycles (besides the two "trivial" fixed points 0 and ∞). In this paper, we prove that they satisfy a weak version of the so-called Ahlfors island property. As a consequence, we can show the density of the repelling periodic points in their Julia sets, and an analogous of Shishikura's result for the forward Julia set J^+ of dissipative Hénon maps.

The class of *Ahlfors island maps* was implicitly present in the work of Epstein [Eps93], see also [RR12, MR12]. Roughly speaking, given an open set $W \subset \mathbb{P}^1$, a holomorphic map $f : W \rightarrow \mathbb{P}^1$ has the N islands property if, given any N Jordan domains with pairwise disjoint closures, one can find univalent inverse branches of f on at least one of these Jordan domains, whose image is close to any given point in the boundary of W (see Definition 2.3 for a precise formulation). A celebrated theorem by Ahlfors [Ahl35] states that every entire or meromorphic map has the 5 islands property (in this case, we have $W = \mathbb{C}$ and the only boundary point is ∞), see also [Ber98]. This remarkable result can be used to give a simple proof of the density of repelling cycles in the Julia set of any transcendental entire or meromorphic map, see for instance [Ber93].

Another class of maps extensively studied by Epstein [Eps93] is the class of *finite type maps*, that is, holomorphic maps with finitely many singular values (see Definition 2.2). Finite type maps with N singular values have the $(N + 1)$ islands property; however, finite type maps form a much smaller class than Ahlfors island maps. For instance, Epstein proved that finite type maps have no wandering or Baker domains and admit only finitely many non-repelling cycles; none of these statements is true in general for Ahlfors island maps. While horn maps of one-variable rational maps are always finite type maps [Eps93], it seems unlikely that this holds true for horn maps of dissipative semi-parabolic Hénon maps in general (indeed, this would be equivalent to proving that only finitely many stable manifolds have tangencies with a certain entire curve Σ , defined below).

In this paper, we introduce the following slightly weaker version of the island property. Observe that in the usual versions, the property below is true for every positive real number instead of just those smaller than $r(z_0)$, see Definition 2.3. On the other hand, here we can specify which points z_0 should be excluded, see Remark 2.4.

Definition 1.1. *Let $W \subset \mathbb{P}^1$ be an open set and $h : W \rightarrow \mathbb{P}^1$ a holomorphic map. We say that h has the small island property if, for every $z_0 \in \mathbb{C}^*$, there exists $r(z_0) > 0$*

such that, for every domain U intersecting ∂W , there exists $\Omega \Subset U \cap W$ such that $h : \Omega \rightarrow \mathbb{D}(z_0, r(z_0))$ is a conformal isomorphism.

Let us emphasize that $r(z_0)$ does not depend on the choice of U , but only on f and z_0 . We will show in Theorem 1.2 that the small island property as in Definition 1.1 is enough to prove the density of the repelling periodic points in the Julia set.

Let now f be a dissipative Hénon map with a semi-parabolic fixed point p of order 2, see Section 3 for the precise definitions. Let \mathcal{B} denote the parabolic basin of the semi-parabolic point, which is a two-dimensional open set with p on its boundary, and Σ the parabolic curve, which can be thought of as the one-dimensional unstable curve of p . By [BSU17], there exist two maps $\phi^t : \mathcal{B} \rightarrow \mathbb{C}$ (usually called the *incoming Fatou coordinate*) and $\psi^o : \mathbb{C} \rightarrow \Sigma$ (usually called the *outgoing Fatou parametrization*) which semi-conjugate the map f on \mathcal{B} and on Σ to the translation by 1 on \mathbb{C} , respectively. The *Hénon-Lavaurs map* (or *transition map*) associated to f is the composition $\mathcal{L}_0 := \psi^o \circ \phi^t$. Observe that \mathcal{L}_0 commutes with f . By [BSU17], \mathcal{L}_0 can be seen as a limit of large iterates of suitable perturbations of f near the semi-parabolic point. We refer to Section 3.4 for more details on this.

By the definition of \mathcal{L}_0 , we immediately see that this map is semi-conjugated to the map $H_f := \phi^t \circ \psi^o : (\psi^o)^{-1}(\mathcal{B}) \rightarrow \mathbb{C}$. This is the *lifted horn map* associated to f . As \mathcal{L}_0 commutes with f , we see that H_f commutes with the translation by 1 on \mathbb{C} . We can then quotient its action by this translation, and obtain a map from a subset of the cylinder \mathbb{C}^* , containing pointed neighbourhoods of 0 and ∞ , to the cylinder itself. By [BSU17], h_f extends to 0 and ∞ . The following is our first main result.

Theorem 1.2. *Let f be a dissipative semi-parabolic Hénon map as above. The horn map h_f has the small island property as in Definition 1.1.*

The following is then a consequence of Theorem 1.2.

Corollary 1.3. *Let f be a dissipative semi-parabolic Hénon map as above. The repelling periodic points of the horn map h_f are dense in its Julia set.*

The proof of Theorem 1.2 is based on techniques from Pesin theory, which were first adapted to this context in [BLS93b], and on local equidistribution results towards the Green currents T^+ and T^- of the Hénon map f , which follow from the local approach to these problems developed in [HOV95, Duj04, DS06, DNS08].

As mentioned above, as an application of Theorem 1.2, we will also deduce an analogous of Shishikura's result for Hénon maps.

Theorem 1.4. *Let f be a dissipative semi-parabolic Hénon map as above. Then, there exists a sequence $f_n \rightarrow f$ of dissipative Hénon maps of the same algebraic degree such that*

$$\dim_H J^+(f_n) \rightarrow 4 \quad \text{as } n \rightarrow \infty.$$

In order to prove Theorem 1.4, we show that every holomorphic map with the small island property can be suitably modified (i.e., can be multiplied by a suitable constant) so that it has arbitrarily large (i.e., close to 2) hyperbolic dimension. Assume for the sake of simplicity that the multiplication is not necessary (this is just a minor technical

point). We can apply this to the horn map h_f , and recall that the Hénon-Lavaurs map \mathcal{L}_0 is the limit of suitable large iterates $f_{\epsilon_n}^n$ of perturbations f_{ϵ_n} of f . By the conjugacy, large hyperbolic sets for h_f give a large limit set for suitable iterates of \mathcal{L}_0 . As these large hyperbolic sets persist under the perturbation $f \mapsto f_{\epsilon_n}$ for n sufficiently large, this leads to hyperbolic limit sets \mathcal{H}_n for suitable iterates of $f_{\epsilon_n}^n$ with large (i.e., close to 2) unstable dimension. As the forward Julia set $J^+(f_\epsilon)$ contains the union of the stable manifolds of the points of \mathcal{H}_n , this leads to the desired result.

Acknowledgments. The authors would like to express their gratitude to Eric Bedford for bringing this problem to their attention. The first author would also like to thank the University of Pisa for the warm welcome and the excellent work conditions.

This project has received funding from the French government through the Programme Investissement d'Avenir (ANR QuaSiDy /ANR-21-CE40-0016, ANR PADAWAN /ANR-21-CE40-0012-01) and from the MIUR Excellence Department Project awarded to the Department of Mathematics, University of Pisa, CUP I57G22000700001. Both authors are part of the PHC Galileo project G24-123.

2. ISLAND PROPERTIES AND CONSEQUENCES

2.1. Ahlfors island maps and finite type maps.

Definition 2.1. *Let X and W be a Riemann surface, with X connected, and $f : W \rightarrow X$ a holomorphic map. The singular value set $S(f)$ of f is the smallest subset of X such that $f : W_0 \setminus f^{-1}(S(f)) \rightarrow X \setminus S(f)$ is a covering map for every connected component W_0 of W .*

As covering maps are surjective, it follows from the above definition that we have $X \setminus f(W) \subset S(f)$.

Definition 2.2. [Eps93] *Let $W \subset \mathbb{P}^1$ be a non-empty open set and $f : W \rightarrow \mathbb{P}^1$ a holomorphic map. We say that f is a finite type map on \mathbb{P}^1 if*

- (1) f is non-constant on every connected component of W ;
- (2) f has no removable singularities;
- (3) $S(f)$ is finite.

Definition 2.3. [RR12, MR12] *Let $W \subset \mathbb{P}^1$ be a non-empty open set and $f : W \rightarrow \mathbb{P}^1$ a holomorphic map. We say that f has the N islands property if, given any N Jordan domains $D_1, \dots, D_N \subset \mathbb{P}^1$ with pairwise disjoint closures and any open set U intersecting ∂W , there exists $1 \leq i_0 \leq N$ and an open set $\Omega \Subset U \cap W$ such that $f : \Omega \rightarrow D_{i_0}$ is a conformal isomorphism. If there exists $N \geq 1$ such that f has the N islands property, we say that f is an Ahlfors island map.*

Remark 2.4. *In Definition 1.1, z_0 must be chosen different from 0 and ∞ . Hence, the small island property as in that definition can be seen as a weaker version of the 3 islands property above. We could give a more general definition of the small N islands property, admitting $N - 1$ exceptions as in the Definition 2.3. The proofs in this section would be the same. We just define the precise version of the property that we will prove for the horn maps of semi-parabolic Hénon maps for simplicity.*

Also note that the 1 island property is vacuously satisfied if $\partial W = \emptyset$, that is, if $f : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is a rational map. This case is however very special, and our arguments will often use the fact that $\partial W \neq \emptyset$.

2.2. Julia sets. We fix in this section an open set $W \subset \mathbb{P}^1$ and a holomorphic map $f : W \rightarrow \mathbb{P}^1$. We give here two natural definitions of the Julia set of f . The first is related to the notion of non-normality. Since $W \neq \mathbb{P}^1$, the definition needs to take into account orbits leaving the domain W .

Definition 2.5 (Definition of the Julia set as non-normality locus). *The Fatou set $F(f)$ of f is the union of all open sets $U \subset \mathbb{P}^1$ such that either*

- (1) $f^n(U) \subset W$ for all $n \in \mathbb{N}$, and $\{f^n : U \rightarrow W\}$ is normal; or
- (2) there exists $n \in \mathbb{N}$ such that $f^n(U) \subset \mathbb{P}^1 \setminus \overline{W}$, where \overline{W} denotes the closure of W in \mathbb{P}^1 .

The set $J_F(f)$ is the complement of $F(f)$ in \mathbb{P}^1 .

Observe that according to this definition, we always have $\partial W \subset J_F(f)$ and $\mathbb{P}^1 \setminus \overline{W} \subset F(f)$. Given a point in the Fatou set, all its orbit is in the Fatou set; conversely, given a point in $J_F(f)$, all preimages are in $J_F(f)$. Moreover, it is clear that repelling periodic points are always in $J_F(f)$. This leads to the second definition, and to the inclusion $J_R(f) \subset J_F(f)$.

Definition 2.6 (Definition of the Julia set by means of repelling periodic points). *The set $J_R(f)$ is the closure of the set of all repelling periodic points of f .*

2.3. A consequence of the small island property. We fix again an open set $W \subset \mathbb{P}^1$ and a holomorphic map $f : W \rightarrow \mathbb{P}^1$. By [Eps93], we have $J_F(f) = J_R(f)$ if f is an Ahlfors island map. In this section we show that the small island property as in Definition 1.1 is still enough to ensure this property. For simplicity, we will denote by W_∞ the interior of $\bigcap_{n \geq 0} f^{-n}(W)$ in the rest of this section.

Lemma 2.7. *Assume that either $W_\infty = \emptyset$ or all connected components of W_∞ are hyperbolic. Then we have $J_F(f) = \overline{\bigcup_{n \geq 0} f^{-n}(\partial W)}$.*

Proof. The inclusion $\overline{\bigcup_{n \geq 0} f^{-n}(\partial W)} \subset J_F(f)$ is always true by the Definition 2.5 of $J_F(f)$.

Conversely, if $W_\infty \neq \emptyset$, W_∞ is completely invariant and $f : W_\infty \rightarrow W_\infty$ is non-increasing for the hyperbolic metric. Therefore, we have $W_\infty \subset F(f)$. Clearly, this inclusion still holds if $W_\infty = \emptyset$.

Let U be a connected open set intersecting $J_F(f)$. By the inclusion proved above, U cannot be contained in W_∞ . Therefore, there exists $n \in \mathbb{N}$ such that $f^n(U) \cap (\mathbb{P}^1 \setminus W) \neq \emptyset$. Moreover, we must have $f^n(U) \cap \partial W \neq \emptyset$, for otherwise we would have $f^n(U) \subset \mathbb{P}^1 \setminus W$. By definition, this would give $U \subset F(f)$, contradicting the assumption $U \cap J_F(f) \neq \emptyset$.

The above proves that, for any open domain U intersecting $J_F(f)$, there exists $n \in \mathbb{N}$ such that $f^{-n}(\partial W) \cap U \neq \emptyset$. Again by definition, we have $f^{-n}(\partial W) \subset J_F(f)$. This shows that $\bigcup_{n \geq 0} f^{-n}(\partial W)$ is indeed dense in $J_F(f)$, and concludes the proof. \square

Theorem 2.8. *Assume that f has the small island property. Then $J_F(f) = J_R(f)$.*

Proof. Recall that the inclusion $J_R(f) \subset J_F(f)$ always holds by definition. Hence, we only have to prove the reversed inclusion.

If W_∞ is non-empty and has a non-hyperbolic component, it is isomorphic to \mathbb{P}^1 , \mathbb{C}^* , or \mathbb{C} . Then, f is either a rational map, a transcendental self-map of \mathbb{C}^* or a transcendental entire map. In all these cases, the result is classical. Therefore, we only need to deal with the case where W_∞ is either empty or has a hyperbolic connected component. In particular, Lemma 2.7 applies.

Fix $z_0 \in J_F(f)$. Since $J_F(f)$ has no isolated points, we may assume without loss of generality that $z_0 \in \mathbb{C}^*$. By Lemma 2.7, we may also assume that $f^n(z_0) \in \partial W$ for some $n \in \mathbb{N}$. Let D be a small disk centered at z_0 small enough so that $f^n : D \rightarrow f^n(D)$ is a branched cover, ramified only possibly at $f^n(z_0)$. We can also assume that the radius of D is smaller than the quantity $r(z_0)$ as in the Definition 1.1 of the small island property. It is enough to show that there exists a repelling point in D .

Let $\Omega \Subset f^n(D)$ be the simply connected domain given by the small island property applied with $U = f^n(D)$. Then, the map $f^{n+1} : f^n(D) \ni \Omega \rightarrow f^n(D)$ is a branched cover with at most one critical value $f^n(z_0)$, which lies outside of $\overline{\Omega}$, and in particular, it is polynomial-like. It follows from, e.g., Douady-Hubbard's Straightening theorem [DH85b] that $f^{n+1} : \Omega \rightarrow f^n(D)$ has a repelling periodic point¹.

Therefore, D contains a repelling periodic point for f . This proves the inclusion $J_F(f) \subset J_R(f)$ and completes the proof. \square

Remark 2.9. A holomorphic family of maps $f_\lambda : W_\lambda \rightarrow \mathbb{P}^1$ is natural [EL92, ABF21] if it is of the form $f_\lambda = \phi_\lambda \circ f \circ \psi_\lambda^{-1}$, where $W \subset \mathbb{P}^1$ is an open set, $f : W \rightarrow \mathbb{P}^1$ a holomorphic map, $W_\lambda := \psi_\lambda(W)$ and $\phi_\lambda, \psi_\lambda : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ are homeomorphisms depending holomorphically on $\lambda \in M$. It is straightforward to check that if f has the small island property, then each map f_λ in a natural family as above also has the small island property. In the proof of Theorem 1.4, we will be interested in the case of a family of maps of the form $h_\lambda = \lambda h$, where h has the small island property and $\lambda \in \mathbb{C}^*$. It is in particular a natural family, with $\phi_\lambda(z) := \lambda z$ and $\psi_\lambda := \text{Id}$.

3. PRELIMINARIES ON HÉNON AND HORIZONTAL-LIKE MAPS

We will consider in this section an automorphism f of \mathbb{C}^2 of the form

$$(3.1) \quad f(z, w) = (p(z) + aw, z),$$

where p is a monic polynomial of degree $d \geq 2$ and a is some constant in \mathbb{C}^* . Any f as above is usually referred to as a (*generalized*) *Hénon map*. Observe that the Jacobian of f is constant, and equal to $|a|$. We say that f is *dissipative* if $|a| < 1$.

By results of Jung [Jun42] and Friedland-Milnor [FM89], every polynomial automorphism f of \mathbb{C}^2 is conjugated (in the group of polynomial automorphisms) to either an *elementary automorphism*, i.e., a map of the form $(z, w) \mapsto (az + p(w), bw + c)$ for some $a, b \in \mathbb{C}^*$, $c \in \mathbb{C}$, and polynomial p of degree $d \geq 0$, or to a *Hénon-type map*, i.e., a finite composition of generalized Hénon maps as in (3.1). As the dynamics of elementary automorphisms is simple to describe, we will just consider in the following maps of the

¹Actually, since this polynomial-like map has only one critical value which is escaping, it can also be proved by elementary means that it has a repelling fixed point, see for instance [Eps93].

second type. For simplicity, we will just consider Hénon maps, but the picture is the same when considering finite compositions as above.

3.1. Filtration property and induced horizontal-like map. Let f be as in (3.1). Friedland and Milnor [FM89] showed that it is possible to decompose \mathbb{C}^2 in a dynamically meaningful way, as follows. Let $\mathbb{D}(0, R) \subset \mathbb{C}$ be the disc of center 0 and radius R . Set

$$\begin{aligned} D_R &:= \mathbb{D}(0, R)^2, \\ V_R^+ &:= \{(z, w) \in \mathbb{C}^2 : |z| > \max(|w|, R)\}, \text{ and} \\ V_R^- &:= \{(z, w) \in \mathbb{C}^2 : |w| > \max(|z|, R)\}. \end{aligned}$$

We also denote by K^+ (resp. K^-) the set of points whose orbit under f (resp. f^{-1}) is bounded.

Lemma 3.1. *The following assertions hold for every R sufficiently large.*

- (1) $f(V_R^+) \subset V_R^+$ and $V_R^+ \cap K^+ = \emptyset$;
- (2) $f(D_R \cup V_R^+) \subset D_R \cup V_R^+$;

Similar assertions hold replacing f , V_R^+ , and K^+ with f^{-1} , V_R^- , and K^- , respectively.

Although our maps will always be globally defined, in the following we will sometimes need to work in a semi-local setting, that we now describe. A *vertical subset* of D_R is a subset of D_R whose closure in \mathbb{C}^2 is disjoint from the *vertical boundary* $\partial\mathbb{D}(0, R) \times \mathbb{D}(0, R)$ of D_R . Similarly, a *horizontal subset* of D_R is a subset of D_R whose closure is disjoint from the *horizontal boundary* $\mathbb{D}(0, R) \times \partial\mathbb{D}(0, R)$.

We fix R sufficiently large for Lemma 3.1 to hold. The map f can be seen as a map from the set $D_R \cap f^{-1}(D_R)$ to the set $D_R \cap f(D_R)$. In particular, since $f^{-1}(D_R) \Subset D_R \cup V^-$ and $f(D_R) \Subset D_R \cup V^+$, f is a holomorphic map from a vertical subset of D_R to a horizontal one and it is a so-called *horizontal-like map*. We refer to [HOV95, Duj04, DS06, DNS08] for the precise definition and their properties. We will use the notation \tilde{f} for the horizontal-like map associated to f as above, when we will need to emphasize it.

3.2. The Green currents T^+ and T^- . Let f be a Hénon map. Denote by G^\pm the functions $G^\pm(z, w) := \lim_{n \rightarrow \infty} d^{-n} \log^+ \|f^{\pm 1}(z, w)\|$, where $\log^+(\cdot) = \max(0, \log(\cdot))$. Such functions, usually called the *Green functions* of f and f^{-1} respectively, are well defined (as the convergences are uniform on every compact subset of \mathbb{C}^2). They are Hölder continuous and plurisubharmonic [FS92, Hub86]. Hence, the $(1, 1)$ -currents given by $T^\pm := dd^c G^\pm$ are positive closed. They are the *Green currents* of f and f^{-1} and they describe – in a quantified sense – the distribution of the iterate of curves under forward and backward iteration of f [BS91a, FS92]. Their support is equal to $J^\pm := \partial K^\pm$, respectively. More precisely, they are the unique positive dd^c -closed currents supported on K^\pm , respectively [DS14]. By [BS91b], the convergence

$$(3.2) \quad d^{-n}[f^{-n}(M)] \rightarrow c_M T^+$$

holds for every locally closed submanifold $M \subset \mathbb{C}^2$ satisfying $M \subset J^+$ or $M \subset X$, where X is algebraic. Here c_M is a constant depending on M , and we have $c_M > 0$ if, for instance, one has $T^-|_M > 0$. A similar property holds for T^- .

In the semi-local setting described in Section 3.1, the convergence above can be improved. For a given large R , recall that we denote by \tilde{f} the associated horizontal-like map on D_R . By [Duj04], we can associate to \tilde{f} its Green current \tilde{T}^+ , which is vertical, and the Green current of \tilde{f}^{-1} , \tilde{T}^- , which is horizontal in D_R . By [Duj04, DNS08], we have

$$(3.3) \quad d^{-n} \tilde{f}^* R \rightarrow c_R \tilde{T}^+$$

for every positive closed vertical current on D_R . Here $c_R > 0$ is a constant depending on R (and it is equal to its *vertical mass*, see [Duj04, DS06]; when R is smooth, it is equal to the mass of the restriction of R to any horizontal line in D_R). It follows from (3.2) and (3.3) that $\tilde{T}^\pm = T_{|D_R}^\pm$.

3.3. The equilibrium measure and Pesin boxes. As the Green functions G^\pm are continuous, the intersection $\mu := T^+ \wedge T^-$ is a well-defined probability measure, and is the unique measure of maximal entropy of f [BLS93b, Sib99]. It satisfies remarkable ergodic properties [BLS93b, BLS93a, Din05, BD24]. We recall here those that we will need in the sequel. We denote by λ^+ and λ^- the two Lyapunov exponents of μ . Recall [BS98] that we have $\lambda^+ > 0$ and $\lambda^- < 0$, hence μ is hyperbolic in the sense of Pesin theory.

By Oseledec's ergodic theorem, there exist a full measure subset \mathcal{R} of the support of μ and two measurable distributions of 1-dimensional subspaces $E^s, E^u: \mathcal{R} \rightarrow T\mathbb{C}^2$ with $E^s(x), E^u(x) \in T_x\mathbb{C}^2$ with the property that for every $x \in \mathcal{R}$, we have $E^s(x) \neq E^u(x)$, $Df(E^{s/u}(x)) = E^{s/u}(f(x))$ and

$$\lim_{n \rightarrow \infty} \|Df^n(v)\| = \lambda^+ \quad \forall v \notin E^s(x) \quad \text{and} \quad \lim_{n \rightarrow \infty} \|Df^{-n}(v)\| = \lambda^- \quad \forall v \notin E^u(x).$$

The angle between E^s and E^u along an orbit is also controlled.

Given $r > 0$ and $x \in \mathcal{R}$, we denote by $B_r^s(x)$ and $B_r^u(x)$ the 1-dimensional affine discs in \mathbb{C}^2 centred at x , of radius r , and whose tangent at x is given by $E^s(x)$ and $E^u(x)$, respectively. By Pesin theory, for every x there is an $r(x)$ such that the stable and unstable manifolds $W^s(x)$ and $W^u(x)$ of x are locally graphs over $B_r^s(x)$ and $B_r^u(x)$, respectively. We denote by $W_r^s(x)$ and $W_r^u(x)$ these local stable and unstable manifolds, respectively.

Fix $r > 0$ and denote by \mathcal{R}_r the set of points $x \in \mathcal{R}$ such that $r(x) \geq r$ ². Let F be a compact subset of \mathcal{R}_r , and assume that the diameter of F is $\ll r$. We denote by $W_r^s(F)$ and $W_r^u(F)$ the union of the sets $W_r^s(x)$ and $W_r^u(x)$ for $x \in F$. We call the set $P := W_r^s(F) \cap W_r^u(F)$ the *Pesin box* [Pes77] generated by F . By [BLS93b] (see also [Duj04]), there exist a compact set P^s which is homeomorphic to $W_r^u(x) \cap W_r^s(F)$ for all $x \in F$ and a compact set P^u which is homeomorphic to $W_r^s(x) \cap W_r^u(F)$ for all $x \in F$. Then, P is homeomorphic to $P^s \times P^u$. Moreover, there exists a neighbourhood $N = N(P)$ of P , biholomorphic to a bidisk, such that (the image of) every $W_N^s(x) := W_r^s(x) \cap N$ (resp. $W_N^u(x) := W_r^u(x) \cap N$) is a vertical (resp. horizontal) graph, and for every $x, y \in P$ the unique intersection point between $W_N^s(x)$ and $W_N^u(x)$ is in P .

It follows from Pesin theory [Pes77] that, up to a negligible set, the support of μ can be covered by means of just countably many Pesin boxes.

²Further conditions should be imposed on \mathcal{R}_r , see [BLS93b, (4.2), (4.3), and (4.4)]. Since we will not need them and they are always true up to a zero-measure subset, we will not focus on this issue here.

3.4. Semi-parabolic dynamics and horn maps. Let us assume from now on that a Hénon map f has a semi-parabolic point of order 2 at the origin \mathbb{O} of \mathbb{C}^2 . A description of the local dynamics of f near the semi-parabolic point is given in [Ued86, Ued91, BSU17]. Following [BSU17], we recall here the definitions and results that we will need in the sequel. We refer to [Lav89, Shi00] for the earlier one-dimensional counterparts of these definitions and results.

Up to suitable changes of coordinates, we can assume that the local form of f near \mathbb{O} is given by

$$f(z, w) = (z + z^2 + O(z^3), bw + O(zw))$$

for some $0 < |b| < 1$ (a more precise development is given in [DL15], but we will not need it here).

We denote by \mathcal{B} the *parabolic basin* of \mathbb{O} , i.e., the open set of points x such that $f^n(x) \rightarrow \mathbb{O}$ as $n \rightarrow \infty$. Observe that, as f is invertible, \mathcal{B} is connected. There exists a holomorphic submersion $\phi^t: \mathcal{B} \rightarrow \mathbb{C}$, called the (one dimensional) *incoming Fatou coordinate*, that semi-conjugates the dynamics of f on \mathcal{B} to a translation by 1 on \mathbb{C} ; i.e., we have

$$(3.4) \quad \phi^t(f(x)) = \phi^t(x) + 1 \quad \forall x \in \mathcal{B}.$$

There also exists a second (open) holomorphic map $\phi_2: \mathcal{B} \rightarrow \mathbb{C}$ such that the map $\Phi = (\phi^t, \phi_2): \mathcal{B} \rightarrow \mathbb{C}^2$ is a biholomorphism which satisfies $\Phi(f(x)) = \Phi(x) + (1, 0)$ for every $x \in \mathcal{B}$. Given any $p \in \mathcal{B}$, the fiber $\{q \in \mathcal{B} : \phi^t(p) = \phi^t(q)\}$ is called the *strongly stable manifold* of p , denoted by $W^{ss}(p)$. It is an injectively immersed entire curve in \mathbb{C}^2 , and it is characterized by the following property:

$$q \in W^{ss}(p) \iff \lim_{n \rightarrow \infty} \frac{1}{n} \log d(f^n(p), f^n(q)) = \log |b|.$$

Let us now consider the set of points converging to \mathbb{O} under the iteration of f^{-1} . This set is an f -invariant complex curve $\Sigma \subset \mathbb{C}^2$, with \mathbb{O} on its boundary. The *Fatou parametrization* of Σ is a holomorphic map $\psi^o: \mathbb{C} \rightarrow \mathbb{C}^2$ satisfying

$$(3.5) \quad f(\psi^o(y)) = \psi^o(y + 1) \quad \forall y \in \mathbb{C}.$$

Definition 3.2. *The map $\mathcal{L}_0 := \psi^o \circ \phi^t: \mathcal{B} \cap \Sigma \rightarrow \Sigma$ is the Hénon-Lavaurs, or transition map of f (associated to the semi-parabolic point \mathbb{O}). The map $H_f := \phi^t \circ \psi^o: (\psi^o)^{-1}(\mathcal{B} \cap \Sigma) \rightarrow \mathbb{C}$ is the lifted horn map of f (associated to the semi-parabolic point \mathbb{O}).*

The following properties of H_f directly follow from its definition and the local description of \mathcal{B} and Σ , see [BSU17, DL15]. The last item is a key point in the characterization of bifurcations in [DL15] by means of homoclinic tangencies.

Proposition 3.3. *The following properties hold.*

- (1) *The domain $\psi^o(\mathcal{B} \cap \Sigma)$ contains the set $\{|\Im z| > R\}$ for every R large enough.*
- (2) *For every $z \in (\psi^o)^{-1}(\mathcal{B})$ and every $w \in \mathbb{C}$, we have $H_f(z) = w$ if and only if $\psi^o(z)$ lies in the intersection of Σ and the strongly stable manifold $\{(x, y) \in \mathcal{B} : \phi^t(x, y) = w\}$.*
- (3) *Given $z \in \mathbb{C}$, we have $H'_f(z) = 0$ if and only if the strongly stable manifold $\{(x, y) \in \mathcal{B} : \phi^t(x, y) = w\}$ is tangent to Σ at $\psi^o(x)$.*

The maps \mathcal{L}_0 and H_f are conjugated to each other, so their dynamics are very similar. In particular, one can use the map $H_f : \mathbb{C} \rightarrow \mathbb{C}$ as a model for the map \mathcal{L}_0 , whose domain and image are in \mathbb{C}^2 . It follows from (3.4) and (3.5) that we have

$$f \circ \mathcal{L}_0 = \mathcal{L}_0 \circ f \quad \text{and} \quad H_f \circ \tau_1 = \tau_1 \circ H_f,$$

where we denote by τ_1 the translation by 1 in \mathbb{C} . In particular, H_f induces a map h_f on (a subset of) the cylinder \mathcal{C} obtained by taking the quotient of \mathbb{C} by the \mathbb{Z} -action of τ_1 . By Proposition 3.3 (1), the domain of h_f contains the two extremities of \mathcal{C} . By [BSU17], h_f extends to such extremities, that we can identify with 0 and ∞ in \mathbb{P}^1 . In particular, we can see h_f as a map from an open subset of \mathbb{P}^1 (containing two neighbourhoods of 0 and ∞) to \mathbb{P}^1 .

Definition 3.4. *The map h_f is the horn map of f (associated to the semi-parabolic point \odot).*

The maps \mathcal{L}_0 , H_f , and h_f are deeply related to the so-called (semi-)parabolic implosion phenomenon for the perturbations of the map f , see [BSU17] and [Lav89, Shi00] for their counterparts in one-dimensional parabolic dynamics. While we will not need results in this direction in the proof of Theorem 1.2, we recall here what we will need in the proof of Theorem 1.4.

Observe that the map \mathcal{L}_0 is actually defined on \mathcal{B} . For every $\alpha \in \mathbb{C}$, define the Hénon-Lavaurs maps of phase α $\mathcal{L}_\alpha : \mathcal{B} \rightarrow \mathbb{C}^2$ as

$$\mathcal{L}_\alpha := \psi^o \circ \tau_\alpha \circ \phi^t,$$

where τ_α denotes the translation by $\alpha \in \mathbb{C}$ in \mathbb{C} . Observe that the image of \mathcal{L}_α is contained in Σ , and that the definition of \mathcal{L}_0 is coherent with its previous definition above.

For small ϵ , we consider holomorphic perturbations f_ϵ of f of the form

$$f_\epsilon(z, w) = (z + z^2 + \epsilon^2 + O(z^2), b_\epsilon w + O(zw)),$$

where in particular b_ϵ depends holomorphically in ϵ . Following [BSU17], we say that a sequence $(n_j, \epsilon_j) \subset (\mathbb{N}, \mathbb{R}^+)^{\mathbb{N}}$ is an α -sequence if $\epsilon_j \rightarrow 0$ and $n_j - \pi/\epsilon_j \rightarrow \alpha$. Observe that this condition implies that $n_j \rightarrow \infty$, and prescribes that the convergence $\epsilon_j \rightarrow 0$ happens tangentially to the positive real axis.

Theorem 3.5 (Bedford-Smillie-Ueda [BSU17]). *Let (n_j, ϵ_j) be an α -sequence. Then,*

$$f_{\epsilon_j}^{n_j} \rightarrow \mathcal{L}_\alpha$$

locally uniformly in \mathcal{B} .

4. PROOF OF THEOREM 1.2

We continue to use the notation of the previous sections. Let $R > 0$ be large enough so that the filtration property in Section 3.1 holds for f and R . We will only be concerned with the dynamics of f inside D_R , where we recall that f can be seen as an invertible horizontal-like map. Fix $z_0 \in \mathbb{C}$ and denote by $S(z_0)$ a connected component of $D_R \cap \{\phi^t(x, y) = z_0\}$. We also let D be a small disk in Σ intersecting the boundary of \mathcal{B} . As we

will see, the proof of Theorem 1.2 will essentially consist in finding suitable intersections between preimages of $S(z_0)$ and images of D .

In order to find such intersections, let us fix an arbitrary Pesin box P for μ , see Section 3.3. Recall that we denote by N a given neighbourhood of P , where the stable/unstable manifolds associated to P can be thought as vertical/horizontal, see Section 3.3.

We first consider the preimages of $S(z_0)$. Here is where the semi-local setting of horizontal-like maps will turn out to be useful, as $S(z_0)$ does not a priori satisfy the conditions for the convergence 3.2.

Lemma 4.1. *We have $d^{-n}(f^n)^*[S(z_0)] \rightarrow T^+$ on D_R .*

Proof. Observe that, up to choosing $R > 0$ large enough, we have $S(z_0) \subset K^+$, which implies that $S(z_0) \cap V_R^+ = \emptyset$. Hence, $S(z_0)$ is a vertical analytic set in D_R . Moreover, we have $[S(z_0)] \wedge [L] = 1$ for every horizontal line L in D_R . The assertion is then a consequence of (3.3). \square

Lemma 4.2. *For μ -a.e. $x_1 \in P$, there exists $n_1 \in \mathbb{N}$ such that $f^{-n_1}(S(z_0))$ intersects $W^u(x_1)$ transversally at some $y_1 \in P$.*

Proof. In the case where $S(z_0)$ is replaced by a disc Δ in the stable manifold of a saddle point, the assertion follows from [BLS93b, Lemma 9.1]. Since the proof only uses that Δ satisfies $d^{-n}[f^{-n}\Delta] \rightarrow T^+$, the same proof applies here thanks to Lemma 4.1. \square

For any $r > 0$, let $\mathcal{S}(r)$ denote the connected component of $\{(x, y) \in D_R : \phi^t(x, y) \in \mathbb{D}(z_0, r)\}$ which contains $S(z_0)$. For any $z \in \mathbb{D}(z_0, r)$, we let $S(z)$ denote the connected component of $\{\phi^t(x, y) = z\}$ contained in $\mathcal{S}(r)$.

By Lemma 4.2, there exist $n_1 \in \mathbb{N}$ and $x_1 \in P$ such that $f^{-n_1}(S(z_0))$ intersects transversally $W_N^u(x_1)$ in P . By the description of the Pesin boxes in Section 3.3 and continuity, we get the following result.

Lemma 4.3. *For every $z_0 \in \mathbb{C}$ there exists $r(z_0) > 0$ such that*

- (1) *the connected component U of $\mathcal{S}(r(z_0)) \cap W_N^u(x_1)$ containing y_1 is simply connected and relatively compact in N ; and*
- (2) *for all $z \in \mathbb{D}(z_0, r(z_0))$, $f^{-n_1}(S(z))$ intersects $W_N^u(x_1)$ transversally at a point in U .*

We now prove a result analogous to Lemma 4.2 for a disk in Σ intersecting the boundary of \mathcal{B} . We first need the following analogous of Lemma 4.1.

Lemma 4.4. *Let $D \subset \Sigma$ be a small disk which intersects $\partial\mathcal{B}$. Then*

- (1) *$T^+|_D > 0$, and*
- (2) *$d^{-n}(f^n)_*[D] \rightarrow c_D T^-$, for some positive constant c_D .*

Proof. By assumption, there exists $x \in \partial\mathcal{B} \cap D$, so that $G^+(x) = 0$. Let us prove that there exists $x_1 \in D$ such that $G^+(x_1) > 0$. Assume, by contradiction, that $G^+ \equiv 0$ on D . Then, we have $D \subset K^+ \cap K^-$. In particular, we have $f^n(D) \subset D_R$ for all $n \in \mathbb{N}$. Therefore, $\{f^n : D \rightarrow \mathbb{C}^2\}$ is a normal family. As there is an open subset of D contained in \mathcal{B} , any limit of any subsequence $f_{|D}^{n_k}$ must be constantly equal to the semi-parabolic

point. This implies that $D \subset \mathcal{B}$ and gives a contradiction with the assumption that $D \cap \partial\mathcal{B} \neq \emptyset$.

Therefore, $G^+ : D \rightarrow \mathbb{R}$ is not constantly equal to 0. On the other hand, it is a non-constant subharmonic function on D which admits a local minimum at $x \in D$. It follows by the maximum principle that G^+ cannot be harmonic on D , which means that $dd^c G^+ > 0$ on D . This proves the first assertion.

By assumption, we have $D \subset \Sigma \subset J^-$. Applying (3.2) (with f^{-1} instead of f), by the first item we have $d^{-n}(f^n)_*[D] \rightarrow c_D T^-$, for some $c_D > 0$. The assertion follows. \square

Lemma 4.5. *For μ -a.e. $x_2 \in P$, there exists $n_2 \in \mathbb{N}$ such that $f^{n_2}(D)$ intersects transversally $W_N^s(x_2)$ at some y_2 in N .*

Proof. As in Lemma 4.2, thanks to Lemma 4.4 the proof is the same as that of [BLS93b, Lemma 9.1]. \square

The following lemma gives the desired tranverse intersections between preimages of disks in stable manifolds in \mathcal{B} and images of disks in Σ intersecting the boundary of \mathcal{B} .

Lemma 4.6. *There are two sequences of integers $N_j, M_j \rightarrow +\infty$ such that the following properties hold.*

- (1) $f^{-N_j}(U) \subset N$ for all j ;
- (2) $\text{diam}(f^{-N_j}(U)) \rightarrow 0$ as $j \rightarrow +\infty$;
- (3) for all $z \in \mathbb{D}(z_0, r(z_0))$, the connected component of $f^{-n_1-N_j}(S(z)) \cap N$ intersecting $f^{-N_j}(U)$ is a vertical graph in N ;
- (4) $f^{M_j}(y_2) \in N$ for all j ;
- (5) the connected component of $f^{n_2+M_j}(D) \cap N$ containing $f^{M_j}(y_2)$ is a horizontal graph in N .

In particular, for all $z \in \mathbb{D}(z_0, r(z_0))$, $f^{-n_1-N_j}(S(z))$ intersects $f^{n_2+M_j}(D)$ transversally at a point in N .

Proof. We fix points x_1, x_2 and integers n_1, n_2 as in Lemmas 4.2 and 4.5.

By Poincaré's recurrence theorem, there exists $N_j \rightarrow \infty$ such that $f^{-N_j}(x_1) \in P$. Since $U \subset W_N^u(x_1)$, we have $f^{-N_j}(U) \subset N$. Up to extracting a subsequence, we may assume that $\lim_{j \rightarrow +\infty} f^{-N_j}(x_1) \rightarrow \hat{x}_1 \in P$, and that we have $f_{|U}^{-N_j} \rightarrow \hat{x}_1$ uniformly. This proves (1) and (2).

Let A_j denote the connected component of $f^{-N_j-n_1}(S(r(z_0))) \cap N$ containing U ; by definition, A_j is a union of pieces of strongly stable manifolds $\{\phi^t = z - n_1 - N_j\}$. Let \tilde{S} be one such piece. By the inclination lemma, each \tilde{S} converges in the C^1 topology to $W_N^s(\hat{x}_1)$, which by Pesin theory (see the discussion in Section 3.3) is a horizontal graph in N . This proves (3).

On the other hand, by a symmetric argument, there exists $M_j \rightarrow +\infty$ such that $f^{M_j}(x_2) \in P$, and $f^{M_j}(x_2) \rightarrow \hat{x}_2 \in P$. Since $f^{n_2}(D)$ intersects $W^s(x_2)$ transversally in P , the inclination lemma implies that the connected component of $f^{M_j+n_2}(D) \cap N$ containing $f^{M_j}(y_2)$ converges in the C^1 topology to $W_N^u(\hat{x}_2)$, which is a horizontal graph in N . This proves (4) and (5). \square

End of the proof of Theorem 1.2. Set $W := (\psi^o)^{-1}(\mathcal{B})$. By the definition of h_f and H_f , and in particular by the fact that H_f commutes with the translation τ_1 , it is enough to prove the following property.

- (\star) for every $z_0 \in \mathbb{C}$ there exists $r(z_0) > 0$ such that for any open set \tilde{W} intersecting ∂W there exist $n \in \mathbb{Z}$ and $V \Subset W \cap (\tilde{W} + n)$ such that $H_f : V \rightarrow \mathbb{D}(z_0, r(z_0))$ is a conformal isomorphism.

Fix $z_0 \in \mathbb{C}$ and, without loss of generality, let \tilde{W} be an open ball intersecting ∂W . Set $D := \psi^o(\tilde{W}) \subset \Sigma$, and observe that D satisfies the assumption of Lemma 4.4. Let also $r(z_0)$ and U be as in Lemma 4.3, n_1, n_2 be as in Lemmas 4.2 and 4.5, and let the sequences $\{N_j\}, \{M_j\}$ be as in Lemma 4.6. By Lemma 4.6, for every $z \in \mathbb{D}(z_0, r(z_0))$ and every j , the set $f^{-n_1-N_j-n_2-M_j}(S(z))$ intersects D transversally at a point in $f^{-n_2-M_j}(U)$. By the second item in Lemma 4.6, we can fix j_0 such that the diameter of $f^{-n_1-N_{j_0}}(U)$ is much smaller than the diameter of N . For convenience, we also set $\tilde{n} := n_1 + n_2 + N_{j_0} + M_{j_0}$.

Set $V := \tilde{n} + (\psi^o)^{-1}(f^{-n_2-M_{j_0}-N_{j_0}}(U)) \subset (\psi^o)^{-1}(D) = \tilde{n} + \tilde{W}$. By the choice of j_0 and Lemma 4.6 (4), we have $V \Subset \tilde{n} + \tilde{W}$. For every $v \in V$, let $u = u(v)$ be the point in U given by $u = f^{n_2+M_{j_0}+N_{j_0}}(\psi^o(v - \tilde{n}))$. The map $v \mapsto u(v)$ is a biholomorphism from V to U . Recall also that, by construction, the map $\phi^t \circ f^{n_1}$ is a biholomorphism between U and $\mathbb{D}(z_0, r(z_0))$. Hence, the map $\tilde{H}(\cdot) := \phi^t \circ f^{\tilde{n}} \circ \psi^o(\cdot - \tilde{n})$ is a biholomorphism from $V \Subset \tilde{n} + \tilde{W}$ to $\mathbb{D}(z_0, r(z_0))$. Recalling that $f \circ \psi^o(\cdot) = \psi^o(\cdot + 1)$, we see that $\tilde{H} = H_f$. Hence, H_f is a biholomorphism from $\tilde{n} + V \Subset \tilde{n} + \tilde{W}$ to $\mathbb{D}(z_0, r(z_0))$. This proves (\star), and the assertion follows. \square

Remark 4.7. *Observe that our proof of Theorem 1.2 does not really need that f is a globally defined Hénon map, but it is enough that it is an invertible horizontal-like map. In particular, the map f does not need to be algebraic.*

5. ALMOST MAXIMAL DIMENSION FOR J^+ AND THEOREM 1.4

5.1. Preliminaries and McMullen's result. In order to prove Theorem 1.4, the idea will be to first construct a hyperbolic (repelling) set of large Hausdorff dimension for some horn map, and then to transfer it to some suitable perturbations of the initial semi-parabolic Hénon map. While we could follow more closely Shishikura's arguments [Shi98], we will use a result by McMullen [McM00], see Theorem 5.4 below, to prove that there are small quasiconformal copies of the Mandelbrot set in the parameter space of horn maps. This will allow us to bypass a part of the proof. In this preliminary section, we recall some terminology and McMullen's result.

Definition 5.1. *Let $W \subset \mathbb{P}^1$ be an open set and $f : W \rightarrow \mathbb{P}^1$ a holomorphic map. We say that $z \in \mathbb{P}^1$ is unramified (for f) if the closure of the set $\{x \in W : \exists n \in \mathbb{N}, f^n(x) = z \text{ and } (f^n)'(x) \neq 0\}$ is dense in $J_F(f)$.*

For rational maps, some points may be completely ramified. For instance, it may happen that every preimage of some point is a critical point. However, we will see in Lemma 5.8 that this is not possible for maps with the small island property.

Definition 5.2. *Let M be a complex manifold and $(f_\lambda)_{\lambda \in M} : W \rightarrow \mathbb{P}^1$ a holomorphic family of holomorphic maps. We say that the family $(f_\lambda)_{\lambda \in M}$ has a local Misiurewicz bifurcation of degree $d \geq 2$ at $\lambda_0 \in M$ if:*

- (1) *there exists a marked critical point c_λ of f_λ and $n \in \mathbb{N}$ such that $x_{\lambda_0} := f_{\lambda_0}^n(c_{\lambda_0})$ is a repelling periodic point;*
- (2) *the map $\lambda \mapsto f_\lambda^n(c_\lambda)$ is not constantly equal to x_{λ_0} in a neighborhood of λ_0 ;*
- (3) *c_{λ_0} is unramified;*
- (4) *the local degree of f_λ^n at c_λ is constantly equal to d in a neighborhood of λ_0 .*

Recall that a critical point c_λ is *marked* if it can be followed holomorphically as a function $\lambda \mapsto c_\lambda = c(\lambda)$ of the parameter λ . Such a critical point is *active* at λ_0 if the sequence $f_\lambda^n(c_\lambda)$ is not normal on any neighbourhood of λ_0 . As in [McM00], we remark that the subtle point in the above definition is the fourth request.

We can now recall the main technical results in [McM00]. We will not use the following proposition, but we quote it since proving a version of it in our context will be a main point of our construction.

Proposition 5.3. *Let M be a complex manifold and $(f_\lambda)_{\lambda \in M} : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ a holomorphic family of rational maps. Assume that a marked critical point c_λ is active and unramified at λ_0 . Then, there exists two sequences $M \ni \lambda_n \rightarrow \lambda_0$ and $\mathbb{N} \ni m_n \rightarrow \infty$ such that, for every $n \in \mathbb{N}$, the family $f_\lambda^{m_n}$ has a local Misiurewicz bifurcation at λ_n .*

We denote by \mathbf{M}_d the *degree d Mandelbrot set*, i.e., the connectedness locus of the family $z^d + c$, for $c \in \mathbb{C}$.

The theorem below was proved by McMullen in the context of holomorphic families of rational maps. However, the arguments are purely local (indeed, the proof consists in finding suitable polynomial-like restrictions of the maps f_λ) and the proof carries through verbatim in our more general setting.

Theorem 5.4 (McMullen [McM00]). *If a holomorphic family $(f_\lambda)_{\lambda \in M} : W \rightarrow \mathbb{P}^1$ has a local Misiurewicz bifurcation of degree d at some parameter λ_0 , then there is a sequence of embeddings $\phi_n : \mathbf{M}_d \rightarrow M$ such that for every $c \in \mathbf{M}_d$, we have $\lambda_n := \phi_n(c) \rightarrow \lambda_0$ and, for every $n \in \mathbb{N}$, f_{λ_n} has a polynomial-like restriction which is hybrid-equivalent to $z^d + c$. Moreover, the quasiconformal distortion of these copies tends to 0 as $n \rightarrow +\infty$.*

In practice, McMullen's results give that, whenever – in a family of rational maps – a bifurcation occurs, one can create local Misiurewicz bifurcations, and thanks to them, by means of a renormalization process, a large set of bifurcation parameters. As a consequence, by [Shi98], one can then find maps with large hyperbolic dimension close to any bifurcation parameter. In the next section, we will adapt the above ideas in the setting of maps with the small island property.

5.2. Large hyperbolic sets from the small island property. In this section, we fix an open set $W \subset \mathbb{P}^1$ and a holomorphic map $h : W \rightarrow \mathbb{P}^1$ satisfying the small island property. We also assume that $\partial W \neq \emptyset$ and that h has at least one critical point (both of these properties hold for horn maps of dissipative semi-parabolic Hénon maps; in particular, the existence of a critical point was proved in [DL15]). For every $\lambda \in \mathbb{C}^*$, we denote $h_\lambda := \lambda h$. The following is the main result of this section.

Proposition 5.5. *For every $\epsilon > 0$ there exists $N \in \mathbb{N}$, $\lambda_1 \in \mathbb{C}^*$, simply connected domains $U, U_1, \dots, U_N \subset W \cap \mathbb{C}^*$, and integers m_1, \dots, m_N such that:*

- (1) *for all $1 \leq i < j \leq N$, we have $U_i, U_j \Subset U$ and $\overline{U_i} \cap \overline{U_j} = \emptyset$;*
- (2) *for all $1 \leq i \leq N$, the map $h_{\lambda_1}^{m_i} : U_i \rightarrow U$ is a conformal isomorphism;*
- (3) *if $c := \max_{1 \leq i \leq N} \sup_{z \in U_i} |(h_{\lambda_1}^{m_i})'(z)|$, then*

$$2 - \epsilon \leq \frac{\log N}{\log c}.$$

In particular, it follows from the Bowen formula [Bow08] that the limit set of the Conformal Iterated Function System (CIFS) given by the holomorphic maps $h_{\lambda_1}^{-m_i} : U \rightarrow U_i$ as in the above statement has Hausdorff dimension at least $2 - \epsilon$.

We begin with the following preliminary result, which we will use to replace the non-normality of the critical orbit near a bifurcation parameter in a family of rational maps.

Lemma 5.6 (Shooting Lemma). *Let $(h_\lambda)_{\lambda \in \mathbb{C}^*}$ be the holomorphic family defined by $h_\lambda = \lambda h$ for every $\lambda \in \mathbb{C}^*$. Fix $\lambda_0 \in \mathbb{C}^*$ and let γ_1, γ_2 be two holomorphic maps defined in a neighborhood of λ_0 and such that $h_{\lambda_0}^n \circ \gamma_1(\lambda_0) \in \partial W$ for some $n \in \mathbb{N}$. Then there exists λ' arbitrarily close to λ_0 such that*

$$h_{\lambda'}^{n+1}(\gamma_1(\lambda')) = \gamma_2(\lambda').$$

In the proof of Lemma 5.6 we will need the following consequence of the argument principle.

Lemma 5.7. *Let V be a Jordan domain, and let f, g be holomorphic functions in a neighborhood of \overline{V} . Suppose that $g(\overline{V}) \subset f(V)$ and $g(\partial V) \cap f(\partial V) = \emptyset$. Then there exists $\lambda \in V$ such that $f(\lambda) = g(\lambda)$.*

Proof of Proposition 5.6. For every $\lambda \in \mathbb{C}^*$, set $G(\lambda) := h_\lambda^n(\gamma_1(\lambda))$ and $g(\lambda) := \lambda^{-1}\gamma_2(\lambda)$. Then, the equation $h_\lambda^{n+1}(\gamma_1(\lambda)) = \gamma_2(\lambda)$ may be rewritten as

$$h_\lambda \circ G(\lambda) = g(\lambda).$$

We will apply Lemma 5.7 to the functions $f(\lambda) := h_\lambda \circ G(\lambda)$ and $g(\lambda)$.

Let D be a disk centered at $g(\lambda_0)$ small enough for the small island property of h to apply (i.e., whose radius is smaller than $r(g(\lambda_0))$). Fix also an $\epsilon > 0$ very small compared to the diameter of D , and also small enough so that $G : \mathbb{D}(\lambda_0, \epsilon) \rightarrow G(\mathbb{D}(\lambda_0, \epsilon))$ is a branched cover without critical points besides possibly λ_0 . By assumption, $G(\mathbb{D}(\lambda_0, \epsilon))$ intersects ∂W .

By the small island property of h (which implies the small island property of h_λ for every $\lambda \in \mathbb{C}^*$, see Remark 2.9), there exists $U \Subset G(\mathbb{D}(\lambda_0, \epsilon))$ such that $f_{\lambda_0}(U) = D$. Let V denote a connected component of $G^{-1}(U)$ inside $\mathbb{D}(\lambda_0, \delta)$. By the choice of ϵ , V is a Jordan domain. By construction, we have $f \circ G(U) = D$. On the other hand, $g(U)$ is contained in the disk $\mathbb{D}(g(\lambda_0), C\epsilon^{1/d})$, where d is the local degree of G at λ_0 and C is a positive constant independent of ϵ . Therefore, for ϵ small enough, we have $g(U) \Subset f \circ G(U)$ and $g(\partial U) \cap f \circ G(\partial U) = \emptyset$. It follows from Lemma 5.7 that there exists $\lambda' \in U$ such that $f \circ G(\lambda') = g(\lambda')$. The proof is complete. \square

Proposition 5.3 has an assumption on the non-ramification of the critical point. In our setting, we will be able to get rid of that assumption thanks to the following lemma.

Lemma 5.8. *Every $z \in \mathbb{C}^*$ is unramified for h .*

Proof. Fix $z \in \mathbb{C}^*$ and let $U \Subset W$ be an open set intersecting $J_F(h)$. Define

$$B^-(z) := \{x \in W : \exists n \in \mathbb{N}, h^n(x) = z \text{ and } (h^n)'(x) \neq 0\}.$$

It is enough to show that $U \cap B^-(z) \neq \emptyset$.

As $U \cap J_F(h) \neq \emptyset$, by Montel's lemma there exists $n \in \mathbb{N}^*$ such that $h^n(U) \cap \partial W \neq \emptyset$. We choose the smallest such n . The map $h^n : U \rightarrow h^n(U)$ has only finitely many critical points. On the other hand, by the small island property, there are infinitely many $y \in h^n(U)$ such that $h(y) = z$ and $h'(y) \neq 0$. In particular, we can find one such y such that there exists $x \in U$ with $h^n(x) = y$ and $(h^n)'(x) \neq 0$. As such x belongs to $U \cap B^-(z)$, this concludes the proof. \square

We can now prove the following version of Proposition 5.3.

Proposition 5.9. *Let $W \subset \mathbb{P}^1$ be an open set with $\partial W \neq \emptyset$ and $h : W \rightarrow \mathbb{P}^1$ a holomorphic map satisfying the small island property and with at least one critical point. There exists $\lambda_0 \in \mathbb{C}^*$ such that the family $(h_\lambda := \lambda h)_{\lambda \in \mathbb{C}^*}$ has a local Misiurewicz bifurcation at λ_0 .*

Proof. Take $\lambda_0 \in \mathbb{C}^*$. We allow ourselves to modify λ_0 in what follows. We let $\gamma_1(\lambda)$ be the motion of a critical point near λ_0 , and $\gamma_2(\lambda)$ the motion of a repelling periodic cycle (which exists by Theorem 1.2 and the implicit function theorem). Observe that, as the maps h_λ have the form $h_\lambda = \lambda h$, the map γ_1 is in fact constant; we will therefore simply write γ_1 instead of $\gamma_1(\lambda)$. As we are allowed to modify λ_0 , we can also assume that $\lambda_0 h(\gamma_1) \in \partial W$. Hence, we are in the assumptions of Lemma 5.6. That lemma shows that, up to slightly modifying λ_0 , we have a Misiurewicz relation at λ_0 . As in [McM00], since we are allowed to slightly perturb the starting parameter (and the critical point is automatically unramified by Lemma 5.8), the first three conditions in Definition 5.2 can be achieved from the above construction. Hence, we only have to show that (up to a further small perturbation) we can get the fourth condition in Definition 5.2. Let us recall that, as in [McM00], this property may fail if the local degree of the critical point at λ_0 is larger than at nearby parameters $\lambda' \neq \lambda_0$.

We assume for simplicity that the integer n in Definition 5.2 is equal to 1. Following [McM00], we denote by a_λ the holomorphic motion of the repelling point giving the Misiurewicz relation at λ_0 . We denote by U a small linearization domain for a_λ , for all λ in given small neighbourhood of λ_0 . We will always work with λ in this neighbourhood. We can also assume that the local degree of c_λ is constant outside of λ_0 . We let b_λ be the motion of a second repelling point, which stays in U for all λ in consideration. By the small island property, there exist preimages of b_λ accumulating on a_λ . We denote by b'_λ one of these preimages. Applying again Lemma 5.6, we can find λ' close to λ_0 such that $f(c_{\lambda'}) = b'_{\lambda'}$. The local degree of f is constant near λ' , and since there are no critical points in U , we see that the same is true for the iterate of f mapping $c_{\lambda'}$ to $b'_{\lambda'}$. The proof is complete. \square

Proof of Proposition 5.5. Fix $\epsilon > 0$ and let $d \geq 2$ be the local degree at a given critical point of h . For any $c \in \mathbb{C}$, denote $g_c(z) := z^d + c$. By [Shi98], there exists $c \in \mathbf{M}_d$, $N \in \mathbb{N}$, simply connected domains $V, V_1, \dots, V_N \subset \mathbb{C}$, and integers m_1, \dots, m_N such that:

- (1) for all $1 \leq i < j \leq N$, we have $V_i, V_j \Subset V$ and $\overline{V_i} \cap \overline{V_j} = \emptyset$;
- (2) for all $1 \leq i \leq N$, $g_c^{m_i} : V_i \rightarrow V$ is a conformal isomorphism;
- (3) if $A := \max_{1 \leq i \leq N} \sup_{z \in V_i} |(g_c^{m_i})'(z)|$, then

$$2 - \frac{\epsilon}{2} \leq \frac{\log N}{\log A}.$$

By Proposition 5.9, there exists a local Misiurewicz bifurcation of degree $d \geq 2$ at some $\lambda_0 \in \mathbb{C}^*$ in the family $(h_\lambda := \lambda h)_{\lambda \in \mathbb{C}^*}$. By Theorem 5.4, there exists a sequence $\lambda_n \in \mathbb{C}^*$ and, for every n , a K_n -quasiconformal homeomorphisms $\phi_n : V \rightarrow \phi_n(V)$ with $K_n \rightarrow 1$ as $n \rightarrow \infty$ which conjugates g_c to a polynomial-like restriction of h_{λ_n} . Then, for n large enough, the open sets $U := \phi_n(V)$ and $U_i := \phi_n(V_i)$ satisfy the conditions in the statement. \square

5.3. Proof of Theorem 1.4. By Theorem 1.2, we can apply Proposition 5.5 to the horn map h_f associated to the semi-parabolic fixed point of f as in Definition 3.4. The following proposition is a rewriting of that statement in terms of the Hénon-Lavaurs maps \mathcal{L}_α . Observe that the multiplicative constant in Proposition 5.5 translates to the phase α in the statement below.

Proposition 5.10. *For every $\epsilon > 0$ there exists $\alpha \in \mathbb{C}$, disjoint open sets $U, U_1, \dots, U_N \subset \Sigma$, and integers m_1, \dots, m_N such that*

- (1) for every $1 \leq i < j \leq N$, we have $U_i, U_j \Subset U$ (in the topology induced by Σ) and $\overline{U_i} \cap \overline{U_j} = \emptyset$;
- (2) for every $1 \leq i \leq N$, the map $\mathcal{L}_\alpha^{m_i} : U_i \rightarrow U$ is a conformal isomorphism;
- (3) if $c := \max_{1 \leq i \leq N} \sup_{z \in U_i} |(\mathcal{L}_\alpha^{m_i})'(z)|$, then

$$(5.1) \quad 2 - \epsilon \leq \frac{\log N}{\log c}.$$

In the proposition above, up to taking preimages, we may assume that U and all the U_i 's have small diameter, and that they are transverse to the strong stable foliation in \mathcal{B} . In particular, we may choose a small open set $V \subset \mathbb{C}^2$ with $V \cap \Sigma = U$ and a coordinate system $(x, y) \in \mathbb{D}^2$ for V in which $U = \mathbb{D} \times \{0\}$ and the strongly stable foliation is the vertical foliation given by $x = c$, for $c \in \mathbb{D}$. We allow ourselves to reduce the size of V (in the transversal direction to Σ) in the following, as well as U and the U_i 's. We only work on V (resp. \mathbb{D}^2) in the following. By a slight abuse of notation, we will still denote by $U_i \times \{0\}$ the images of the U_i in the chart (x, y) . Moreover, given two vertical subsets $A, B \subset \mathbb{D} \times \mathbb{D}$, we will write $A \Subset B$ whenever $A \cap \{y = y_0\} \Subset B \cap \{y = y_0\}$ for every $y_0 \in \mathbb{D}$. For every $1 \leq i \leq N$, we also denote $V_i := U_i \times \mathbb{D}$ and by L_i the reading of $\mathcal{L}_\alpha^{m_i}$ in the coordinates $(x, y) \in \mathbb{D}^2$. We also denote by L the holomorphic map on $\cup V_i$ which is equal to L_i on V_i . Observe that, in particular, with the above notations we have

$$(5.2) \quad L_i^{-1}(V_j) \Subset V_i \text{ for all } i, j \quad \text{and} \quad L^{-1}(\cup V_i) \Subset \cup V_i.$$

By Theorem 3.5, given an α -sequence (ϵ_j, n_j) , we have $f_{\epsilon_j}^{n_j} \rightarrow \mathcal{L}_\alpha$ locally uniformly in \mathcal{B} . For simplicity of notation, we will fix an α -sequence of the form (ϵ_n, n) , so that the convergence takes the form

$$(5.3) \quad f_{\epsilon_n}^n \rightarrow \mathcal{L}_\alpha.$$

For every n and i , we denote by $g_{n,i}$ the reading in the coordinates $(x, y) \in \mathbb{D}^2$ of the restriction of $f_{\epsilon_n}^{nm_i}$ to V_i . We also denote by g_n the holomorphic map which is equal to $g_{n,i}$ on V_i . Then, in the coordinates (x, y) , for every n sufficiently large, g_n can be seen as a horizontal-like map [DNS08] from a vertical subset to a horizontal subset of $\mathbb{D} \times \mathbb{D}$. Up to slightly reducing the U_i and V , this vertical subset is actually a subset of $\cup V_i$. The horizontal subset consists of N small horizontal sets, contained in a small neighbourhood of $U = \{0\} \times \mathbb{D}$.

It follows from (5.2) and (5.3) that, for every n sufficiently large and up to slightly reducing the U_i 's, we have

$$(5.4) \quad g_{n,i}^{-1}(V_j) \Subset V_i \text{ for all } i, j \quad \text{and} \quad g_n^{-1}(\cup V_i) \Subset \cup V_i.$$

Let us denote by K_n^+ the set of points in \mathbb{D}^2 whose orbit under g_n never leaves \mathbb{D}^2 (which corresponds to point never leaving V under appropriate iterates of $f_{\epsilon_n}^n$). It follows from (5.4) that, for all n sufficiently large, K_n^+ is a collection of vertical graphs in $\mathbb{D} \times \mathbb{D}$, which are stable manifolds in \mathbb{D}^2 . Similarly, we can also consider the horizontal set K_n^- which is a union of unstable manifolds in \mathbb{D}^2 . The intersection $C_n := K_n^+ \cap K_n^-$ is a Cantor set, on which the action of $f_{\epsilon_n}^n$ is uniformly hyperbolic. We denote by μ_n the measure of maximal entropy $\log N$ on C_n . This measure admits two Lyapunov exponents $\chi_n^- < 0 < \chi_n^+$. The exponent χ_n^+ can be estimated by means of the transversal contraction of the vertical sets V_i under the inverse iteration of g_n . In particular, we deduce the following upper bound from (5.3) and the definition of c in Proposition 5.10:

Lemma 5.11. *Fix $\delta > 1$ and let c be as in Proposition 5.10. Then, for all $n \in \mathbb{N}$ large enough, we have*

$$\chi_n^+ < \delta \log c.$$

Let $W^u(p)$ denote a generic unstable manifold of some $p \in C_n$, and let ν_n denote the conditional measure of μ_n on $W^u(p)$. It follows from [LY85] that the Hausdorff dimension of ν_n is equal to $(\log N)/\chi_n^+$. As a consequence, we deduce from Lemma 5.11 that, for every $\delta > 1$, the Hausdorff dimension of ν_n is larger than $\delta^{-1} \log N/c$ for every n sufficiently large.

Recall that through every $p \in C_n$ there is a stable manifold $W^s(p)$ which is a vertical graph in $\mathbb{D} \times \mathbb{D}$ and an unstable manifold $W^u(p)$ which is a horizontal graph in $\mathbb{D} \times \mathbb{D}$; in particular, $W^s(p)$ intersects every horizontal graph at exactly one point. In particular, for every $y_0 \in \mathbb{D}$ there is a well-defined holonomy map $\phi_{y_0} : \text{supp } \nu_n \rightarrow \mathbb{D}$, associated to the stable foliation of g_n in \mathbb{D}^2 , between the transversals $W^u(p)$ and the horizontal disk $\{y = y_0\}$. By [Lyu99] (see also [LP21]), the map ϕ_{y_0} is Lipschitz continuous. It follows that $K_n^+ = \bigcup_{y \in \mathbb{D}} \phi_y(\text{supp } \nu_n)$ has Hausdorff dimension at least $2 + \delta^{-1}(2 - \epsilon)$. By the relation (5.1) between N and c , this implies that we have

$$\dim_H(K_n^+) \geq 2 + \delta^{-1}(2 - \epsilon)$$

for every n sufficiently large. Up to taking δ sufficiently close to 1 and n sufficiently large, we deduce the lower bound $\dim_H(K_n^+) \geq 4 - 2\epsilon$. As the local coordinates (x, y) are conformal and the image of K_n^+ is contained in $J^+(f_{\epsilon_n}^n) = J^+(f_{\epsilon_n})$, this concludes the proof of Theorem 1.4.

REFERENCES

- [Aba01] Marco Abate. The residual index and the dynamics of holomorphic maps tangent to the identity. *Duke Math. J.*, 107(1):173–207, 2001.
- [Aba05] Marco Abate. Holomorphic classification of 2-dimensional quadratic maps tangent to the identity. *Surikaisekikenkyusho Kokyuroku*, (1447):1–14, 2005.
- [ABD⁺16] Matthieu Astorg, Xavier Buff, Romain Dujardin, Han Peters, and Jasmin Raissy. A two-dimensional polynomial mapping with a wandering Fatou component. *Ann. of Math. (2)*, 184(1):263–313, 2016.
- [ABF21] Matthieu Astorg, Anna Miriam Benini, and Núria Fagella. Bifurcation loci of families of finite type meromorphic maps. *arXiv preprint arXiv:2107.02663*, 2021.
- [ABTP23] Matthieu Astorg, Luka Boc Thaler, and Han Peters. Wandering domains arising from Lavaurs maps with Siegel disks. *Anal. PDE*, 16(1):35–88, 2023.
- [Ahl35] Lars Ahlfors. Zur Theorie der Überlagerungsflächen. *Acta Math.*, 65(1):157–194, 1935.
- [BC12] Xavier Buff and Arnaud Chéritat. Quadratic Julia sets with positive area. *Ann. of Math. (2)*, 176(2):673–746, 2012.
- [BD24] Fabrizio Bianchi and Tien-Cuong Dinh. Every complex Hénon map is exponentially mixing of all orders and satisfies the CLT. *Forum Math. Sigma*, 12:Paper No. e4, 12, 2024.
- [Ber93] Walter Bergweiler. Iteration of meromorphic functions. *Bull. Amer. Math. Soc. (N.S.)*, 29(2):151–188, 1993.
- [Ber98] Walter Bergweiler. A new proof of the Ahlfors five islands theorem. *J. Anal. Math.*, 76:337–347, 1998.
- [Bia19] Fabrizio Bianchi. Parabolic implosion for endomorphisms of \mathbb{C}^2 . *J. Eur. Math. Soc. (JEMS)*, 21(12):3709–3737, 2019.
- [BLS93a] Eric Bedford, Mikhail Lyubich, and John Smillie. Distribution of periodic points of polynomial diffeomorphisms of \mathbb{C}^2 . *Invent. Math.*, 114(2):277–288, 1993.
- [BLS93b] Eric Bedford, Mikhail Lyubich, and John Smillie. Polynomial diffeomorphisms of \mathbb{C}^2 . IV. The measure of maximal entropy and laminar currents. *Invent. Math.*, 112(1):77–125, 1993.
- [Bow08] Rufus Bowen. *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, volume 470 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, revised edition, 2008. With a preface by David Ruelle.
- [BS91a] Eric Bedford and John Smillie. Polynomial diffeomorphisms of \mathbb{C}^2 : currents, equilibrium measure and hyperbolicity. *Invent. Math.*, 103(1):69–99, 1991.
- [BS91b] Eric Bedford and John Smillie. Polynomial diffeomorphisms of \mathbb{C}^2 . II. Stable manifolds and recurrence. *J. Amer. Math. Soc.*, 4(4):657–679, 1991.
- [BS98] Eric Bedford and John Smillie. Polynomial diffeomorphisms of \mathbb{C}^2 . V. Critical points and Lyapunov exponents. *J. Geom. Anal.*, 8(3):349–383, 1998.
- [BSU17] Eric Bedford, John Smillie, and Tetsuo Ueda. Semi-parabolic bifurcations in complex dimension two. *Comm. Math. Phys.*, 350(1):1–29, 2017.
- [CS15] Davoud Cheraghi and Mitsuhiro Shishikura. Satellite renormalization of quadratic polynomials. *arXiv preprint arXiv:1509.07843*, 2015.
- [DH84] Adrien Douady and John H. Hubbard. *Étude dynamique des polynômes complexes. Partie I*, volume 84-2 of *Publications Mathématiques d’Orsay [Mathematical Publications of Orsay]*. Université de Paris-Sud, Département de Mathématiques, Orsay, 1984.
- [DH85a] Adrien Douady and John H. Hubbard. *Étude dynamique des polynômes complexes. Partie II*, volume 85-4 of *Publications Mathématiques d’Orsay [Mathematical Publications of Orsay]*. Université de Paris-Sud, Département de Mathématiques, Orsay, 1985. With the collaboration of P. Lavaurs, Tan Lei and P. Sentenac.
- [DH85b] Adrien Douady and John H. Hubbard. On the dynamics of polynomial-like mappings. *Ann. Sci. École Norm. Sup. (4)*, 18(2):287–343, 1985.
- [Din05] Tien-Cuong Dinh. Decay of correlations for Hénon maps. *Acta Math.*, 195:253–264, 2005.
- [DL15] Romain Dujardin and Mikhail Lyubich. Stability and bifurcations for dissipative polynomial automorphisms of \mathbb{C}^2 . *Invent. Math.*, 200(2):439–511, 2015.

- [DNS08] Tien-Cuong Dinh, Viêt-Anh Nguyễn, and Nessim Sibony. Dynamics of horizontal-like maps in higher dimension. *Adv. Math.*, 219(5):1689–1721, 2008.
- [Dou94] Adrien Douady. Does a Julia set depend continuously on the polynomial? In *Complex dynamical systems (Cincinnati, OH, 1994)*, volume 49 of *Proc. Sympos. Appl. Math.*, pages 91–138. Amer. Math. Soc., Providence, RI, 1994.
- [DS06] Tien-Cuong Dinh and Nessim Sibony. Geometry of currents, intersection theory and dynamics of horizontal-like maps. *Ann. Inst. Fourier (Grenoble)*, 56(2):423–457, 2006.
- [DS14] Tien-Cuong Dinh and Nessim Sibony. Rigidity of Julia sets for Hénon type maps. *J. Mod. Dyn.*, 8(3-4):499–548, 2014.
- [DSZ97] Adrien Douady, Pierrette Sentenac, and Michel Zinsmeister. Implosion parabolique et dimension de Hausdorff. *C. R. Acad. Sci. Paris Sér. I Math.*, 325(7):765–772, 1997.
- [Duj04] Romain Dujardin. Hénon-like mappings in \mathbb{C}^2 . *Amer. J. Math.*, 126(2):439–472, 2004.
- [Éca85] Jean Écalle. *Les fonctions résurgentes. Tome III*, volume 85-5 of *Publications Mathématiques d’Orsay [Mathematical Publications of Orsay]*. Université de Paris-Sud, Département de Mathématiques, Orsay, 1985. L’équation du pont et la classification analytique des objets locaux. [The bridge equation and analytic classification of local objects].
- [EL92] Alexandre È. Erëmenko and Mikhail Lyubich. Dynamical properties of some classes of entire functions. *Ann. Inst. Fourier (Grenoble)*, 42(4):989–1020, 1992.
- [Eps93] Adam L. Epstein. *Towers of finite type complex analytic maps*. PhD thesis, City University of New York, 1993.
- [FM89] Shmuel Friedland and John Milnor. Dynamical properties of plane polynomial automorphisms. *Ergodic Theory Dynam. Systems*, 9(1):67–99, 1989.
- [FS92] John Erik Fornæss and Nessim Sibony. Complex Hénon mappings in \mathbb{C}^2 and Fatou-Bieberbach domains. *Duke Math. J.*, 65(2):345–380, 1992.
- [HOV95] John H. Hubbard and Ralph W. Oberste-Vorth. Hénon mappings in the complex domain. II. Projective and inductive limits of polynomials. In *Real and complex dynamical systems (Hillerød, 1993)*, volume 464 of *NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci.*, pages 89–132. Kluwer Acad. Publ., Dordrecht, 1995.
- [Hub86] John H. Hubbard. The Hénon mapping in the complex domain. In *Chaotic dynamics and fractals (Atlanta, Ga., 1985)*, volume 2 of *Notes Rep. Math. Sci. Engrg.*, pages 101–111. Academic Press, Orlando, FL, 1986.
- [IS06] Hiroyuki Inou and Mitsuhiro Shishikura. The renormalization for parabolic fixed points and their perturbation. *preprint*, 2006.
- [Jun42] Heinrich W. E. Jung. Über ganze birationale Transformationen der Ebene. *J. Reine Angew. Math.*, 184:161–174, 1942.
- [Lav89] Pierre Lavaurs. *Systemes dynamiques holomorphes: explosion de points périodiques paraboliques*. PhD thesis, Paris 11, 1989.
- [LP21] Mikhail Lyubich and Han Peters. Structure of partially hyperbolic Hénon maps. *J. Eur. Math. Soc. (JEMS)*, 23(9):3075–3128, 2021.
- [LY85] François Ledrappier and Lai-Sang Young. The metric entropy of diffeomorphisms. II. Relations between entropy, exponents and dimension. *Ann. of Math. (2)*, 122(3):540–574, 1985.
- [Lyu99] Mikhail Lyubich. Feigenbaum-Couillet-Tresser universality and Milnor’s hairiness conjecture. *Ann. of Math. (2)*, 149(2):319–420, 1999.
- [McM00] Curtis T. McMullen. The Mandelbrot set is universal. In *The Mandelbrot set, theme and variations*, volume 274 of *London Math. Soc. Lecture Note Ser.*, pages 1–17. Cambridge Univ. Press, Cambridge, 2000.
- [MR12] Volker Mayer and Lasse Rempe. Rigidity and absence of line fields for meromorphic and Ahlfors islands maps. *Ergodic Theory Dynam. Systems*, 32(5):1691–1710, 2012.
- [Pes77] Jakov B. Pesin. Characteristic Ljapunov exponents, and smooth ergodic theory. *Uspehi Mat. Nauk*, 32(4(196)):55–112, 287, 1977.
- [PV20] Han Peters and Liz Vivas. Parabolic implosion. *Notices Amer. Math. Soc.*, 67(8):1095–1103, 2020.

- [RR12] Lasse Rempe and Philip J. Rippon. Exotic Baker and wandering domains for Ahlfors islands maps. *J. Anal. Math.*, 117:297–319, 2012.
- [Shi98] Mitsuhiro Shishikura. The Hausdorff dimension of the boundary of the Mandelbrot set and Julia sets. *Ann. of Math. (2)*, 147(2):225–267, 1998.
- [Shi00] Mitsuhiro Shishikura. Bifurcation of parabolic fixed points. In *The Mandelbrot set, theme and variations*, volume 274 of *London Math. Soc. Lecture Note Ser.*, pages 325–363. Cambridge Univ. Press, Cambridge, 2000.
- [Sib99] Nessim Sibony. Dynamique des applications rationnelles de \mathbf{P}^k . In *Dynamique et géométrie complexes (Lyon, 1997)*, volume 8 of *Panor. Synthèses*, pages 97–185. Soc. Math. France, Paris, 1999.
- [Tan98] Lei Tan. Hausdorff dimension of subsets of the parameter space for families of rational maps. (A generalization of Shishikura’s result). *Nonlinearity*, 11(2):233–246, 1998.
- [Ued86] Tetsuo Ueda. Local structure of analytic transformations of two complex variables. I. *J. Math. Kyoto Univ.*, 26(2):233–261, 1986.
- [Ued91] Tetsuo Ueda. Local structure of analytic transformations of two complex variables. II. *J. Math. Kyoto Univ.*, 31(3):695–711, 1991.
- [Vor81] S. M. Voronin. Analytic classification of germs of conformal mappings $(\mathbf{C}, 0) \rightarrow (\mathbf{C}, 0)$. *Funktsional. Anal. i Prilozhen.*, 15(1):1–17, 96, 1981.
- [Zin98] Michel Zinsmeister. Fleur de Leau-Fatou et dimension de Hausdorff. *C. R. Acad. Sci. Paris Sér. I Math.*, 326(10):1227–1232, 1998.

UNIVERSITÉ D’ORLÉANS, INSTITUT DENIS POISSON, UMR CNRS 7013, 45067 ORLÉANS CEDEX 2, FRANCE

Email address: `matthieu.astorg@univ-orleans.fr`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127 PISA, ITALY

Email address: `fabrizio.bianchi@unipi.it`